

## Horizon 2020 Work programme

Food Security, Sustainable Agriculture and Forestry, Marine, Maritime and Inland Water Research and the Bioeconomy *Call* H2020-FNR-2020: Food and Natural Resources *Topic name* FNR-16-2020: ENZYMES FOR MORE ENVIRONMENT-FRIENDLY CONSUMER PRODUCTS

## FuturEnzyme:

Technologies of the Future for Low-Cost Enzymes for Environment-Friendly Products Final ID: 101000327

27 December 2022



# STANDARD ASSAYS, ANALYTICS AND CALCULATIONS FOR MONITORING ENZYMATIC PERFORMANCE D3.2

WOLFGANG STREIT UHAM Department of Microbiology and Biotechnology, Ohnhorststrasse 18, 22609, Hamburg, Germany

## Document information sheet

Work package:	WP3, Activity-based bio-prospecting for enzymes
Authors:	UHAM (Wolfgang R. Streit, Jennifer Chow, Pablo Pérez García), CSIC (Manuel Ferrer, Patricia Molina), Bangor (Peter Golyshin, Alexander Yakunin, Olga Golyshina), UDUS (Karl-Erich Jaeger, Stephan Thies, Alexander Bollinger, Rebecka Molitor), CNR (Michael Yakimov), IST-ID (Carla de Carvalho, Pedro Fernandes)
Document version:	2
Date:	27.12.2022
Starting date:	01/06/2021
Duration:	48 months
Lead beneficiary:	UHAM
Participant(s):	UHAM, CSIC, BANGOR, UDUS, IST-ID, CNR
Dissemination level	Confidential, only for members of the consortium (including the Commission Services)
Туре	Report
Due date (months)	2
Contact details:	Wolfgang Streit, wolfgang.streit@uni-hamburg.de; Manuel Ferrer, mferrer@icp.csic.es

# Summary

Abbreviations	5
1. Scope of the Deliverable	6
2. Reasons for the update	6
3. Origin of the Deliverable	6
4. Protocols for DETERGENT APPLICATIONS	6
4.1. pH shift liquid protocols for quantifying esterase-lipase activity	11
4.1.1. Using Phenol Red (pH 8.0) as a pH indicator	11
4.1.2. Using <i>p</i> -nitrophenol (pH 7.0) as a pH indicator	12
4.1.3. Using nitrazine yellow as a pH indicator	12
4.1.4. Using fluorescein as pH indicator	13
4.2. Liquid protocol for quantifying esterase-lipase activity with chromogenic esters	13
4.2.1. Using short-medium $\alpha$ - or $\beta$ -naphthyl and <i>p</i> -nitrophenol esters	13
4.2.2. Using short-large <i>p</i> -nitrophenol esters	13
4.2.3. Using medium-large <i>p</i> -nitrophenol esters	14
4.3. Liquid protocol for quantifying esterase-lipase activity with non-chromogenic esters	14
4.3.1. Using with fatty acid cholesterol esters	14
4.3.2. Using insoluble substrates (waxes and lipids) and pH-stat	14
4.4. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits	15
4.4.1. NEFA-HR(2) kit	15
4.5. Agar plate for screening esterase-lipase with $lpha$ -naphthyl ester	16
4.6. Agar plate tributyrin screening esterase-lipase assay	17
4.7. Agar plate coconut oil screening lipase assay	17
4.8. Agar plate cocoa butter screening lipase assay	17
4.9. Agar plate olive oil lipase screening assay	
4.10. Agar plate palm oil screening lipase assay	18
4.11. Agar plate egg yolk phospholipase assay	19
4.12. Liquid assay for investigating effect of laundry detergents	19
4.13. Protocol for preparing cell suspensions for activity assays	19
5. Protocols relevant for TEXTILE APPLICATIONS	19
5.1. Agar plate polyesterase screening assay with PET-nanoparticles	22
5.1.1. Protocol for the preparation of PET-nanoparticle solution	22
5.1.2. Preparation of PET-nanoparticle screening plates	22
5.2. Agar plate polyesterase screening assay with polylactic acid (PLA) emulsion	22
5.3. Agar plate polyesterase screening assay with PET or PLA polymers	23
5.4. Agar plate polyesterase screening assay with Impranil DLN	23

5.6. Agar plate polyesterase screening assay with polycaprolactone (PCL) nanoparticles       23         5.7. Agar plate protease skim milk screening assay       24         5.8. Agar plate protease casein screening assay       24         5.9. Agar plate cellulase screening assay       24         5.10. Agar plate oxidoreductase (laccase/oxidase) screening assay       25         5.11. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits       25         5.12. Liquid polyesterase quantification assay with polylactic acid (PLA) powder       27         5.13. Liquid polyesterase quantification assay using Impranil DLN-SD.       27         5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC       27         5.15. Liquid oxidoreductase (laccase/oxidase) quantification assay       27         5.16. Liquid amylase quantification assay with chromogenic substrates       28         5.17. Liquid amylase quantification assay with polymeric substrates       28         5.19. Liquid proteolytic activity assays using P-nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using p-nitroanalide chromophores       29         5.21. Liquid and yeability modification of fabrics       29         5.22. Liquid hydrophilic and dyeability modification of fabrics       32         5.23. Liquid anino modification of polyamide fabrics       30<		5.5. Agar plate polyesterase screening assay with polycaprolactone diol (PCDMn530)	23
5.8. Agar plate protease casein screening assay       24         5.9. Agar plate oxidoreductase (laccase/oxidase) screening assay.       24         5.10. Agar plate oxidoreductase (laccase/oxidase) screening assay.       25         5.11. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits       25         5.12. Liquid polyesterase quantification assay with polylactic acid (PLA) powder       27         5.13. Liquid polyesterase quantification assay using Impranil DLN-SD.       27         5.14. Liquid protocol for the quantification assay with polylactic acid (PLA) powder       27         5.15. Liquid oxidoreductase (laccase/oxidase) quantification assay       27         5.16. Liquid amylase quantification assay with chromogenic substrates       28         5.17. Liquid amylase quantification assay with polymeric substrates       28         5.18. Liquid proteolytic activity assays using <i>p</i> -nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores       29         5.21. Liquid anylor polymeric activity assays using <i>a</i> -nitroanalide chromophores       29         5.22. Liquid proteolytic activity assays using <i>a</i> -nitroanalide chromophores       29         5.23. Liquid proteolytic activity assays using <i>a</i> -nitroanalide chromophores       29         5.24. Dye liquid discoloration test       31         6.1. Liquid colorimetric a		5.6. Agar plate polyesterase screening assay with polycaprolactone (PCL) nanoparticles	. 23
5.9. Agar plate cellulase screening assay       24         5.10. Agar plate oxidoreductase (laccase/oxidase) screening assay.       25         5.11. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits       25         5.12. Liquid polyesterase quantification assay with polylactic acid (PLA) powder       27         5.13. Liquid polyesterase quantification assay using Impranil DLN-SD       27         5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC       27         5.15. Liquid oxidoreductase (laccase/oxidase) quantification assay       27         5.16. Liquid anylase quantification assay with chromogenic substrates       28         5.17. Liquid anylase quantification assay with polymeric substrates       28         5.18. Liquid proteolytic activity assays using P-nitroanalide chromophores       29         5.19. Liquid proteolytic (alcalase) activity assays using <i>p</i> -nitroanalide chromophores       29         5.20. Liquid proteolytic activity assays using azocasein       29         5.21. Liquid annification of polyamide fabrics       30         5.22. Liquid and discolaration test       31         6.1 Protocols relevant for COSMETIC APPLICATIONS       32         6.1. Liquid colorimetric assay for hyaluronidase activity       33         6.1.2 BSA method       33         6.1.3 BCA method       34 <t< td=""><td></td><td>5.7. Agar plate protease skim milk screening assay</td><td> 24</td></t<>		5.7. Agar plate protease skim milk screening assay	24
5.10. Agar plate oxidoreductase (laccase/oxidase) screening assay.       25         5.11. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits       25         5.12. Liquid polyesterase quantification assay with polylactic acid (PLA) powder       27         5.13. Liquid polyesterase quantification assay using Impranil DLN-SD.       27         5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC.       27         5.15. Liquid oxidoreductase (laccase/oxidase) quantification assay       27         5.16. Liquid amylase quantification assay with chromogenic substrates       28         5.17. Liquid amylase quantification assay using FITC-labelled protein       28         5.19. Liquid proteolytic activity assays using <i>p</i> -nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores       29         5.21. Liquid proteolytic activity assays using azocasein       29         5.22. Liquid hydrophilic and dyeability modification of fabrics       29         5.23. Liquid amino modification test       31         6.10. DNS method       33         6.11. DNS method       33         6.12. BSA method       33         6.13. BCA method       34         6.14. DMAB method       34         6.14. daalysis of hyaluronic acid oligosaccharides by HPAE		5.8. Agar plate protease casein screening assay	24
5.11. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits       25         5.11. NEFA-HR(2) kit       25         5.12. Liquid polyesterase quantification assay with polylactic acid (PLA) powder       27         5.13. Liquid polyesterase quantification assay using Impranil DLN-SD       27         5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC       27         5.15. Liquid oxidoreductase (laccase/oxidase) quantification assay       27         5.16. Liquid amylase quantification assay with chromogenic substrates       28         5.17. Liquid amylase quantification assay using Proceed protein       28         5.18. Liquid proteolytic activity assays using FITC-labelled protein       28         5.19. Liquid proteolytic (alcalase) activity assays using <i>p</i> -nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores       29         5.21. Liquid aproteolytic activity assays using azocasein       29         5.22. Liquid hydrophilic and dyeability modification of fabrics       29         5.23. Liquid amino modification of polyamide fabrics       30         5.24. Dye liquid discoloration test       31         6.11. DNS method       33         6.1.2. BSA method       33         6.1.3. BCA method       34         6.1.4. DMAB		5.9. Agar plate cellulase screening assay	24
5.11.1. NEFA-HR(2) kit       25         5.12. Liquid polyesterase quantification assay with polylactic acid (PLA) powder       27         5.13. Liquid polyesterase quantification assay using Impranil DLN-SD       27         5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC       27         5.15. Liquid anylase quantification assay with chromogenic substrates       28         5.17. Liquid anylase quantification assay with polymeric substrates       28         5.18. Liquid proteolytic activity assays using FITC-labelled protein       28         5.19. Liquid proteolytic (alcalase) activity assays using <i>p</i> -nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores       29         5.21. Liquid anylophilic and dyeability modification of fabrics       29         5.22. Liquid hydrophilic and dyeability modification of fabrics       29         5.23. Liquid amino modification test       31         6. Protocols relevant for COSMETIC APPLICATIONS       32         6.1. Liquid colorimetric assay for hyaluronidase activity       33         6.1.2. BSA method       33         6.1.4. DMAB method       34         6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD       34         6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD       34      <		5.10. Agar plate oxidoreductase (laccase/oxidase) screening assay	. 25
5.12. Liquid polyesterase quantification assay with polylactic acid (PLA) powder       27         5.13. Liquid polyesterase quantification assay using Impranil DLN-SD       27         5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC       27         5.15. Liquid avidoreductase (laccase/oxidase) quantification assay       27         5.16. Liquid amylase quantification assay with chromogenic substrates       28         5.17. Liquid amylase quantification assay with polymeric substrates       28         5.18. Liquid proteolytic activity assays using FITC-labelled protein       28         5.19. Liquid proteolytic (alcalase) activity assays using p-nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using p-nitroanalide chromophores       29         5.21. Liquid proteolytic activity assays using azocasein       29         5.22. Liquid hydrophilic and dyeability modification of fabrics       29         5.23. Liquid amino modification of polyamide fabrics       30         5.24. Dye liquid discoloration test       31         6.1. Liquid colorimetric assay for hyaluronidase activity       32         6.1.1. DNS method       33         6.1.2. BSA method       34         6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD       34         6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD <td></td> <td>5.11. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits</td> <td> 25</td>		5.11. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits	25
5.13. Liquid polyesterase quantification assay using Impranil DLN-SD.       27         5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC.       27         5.15. Liquid anylase quantification assay with chromogenic substrates       28         5.17. Liquid anylase quantification assay with polymeric substrates       28         5.18. Liquid proteolytic activity assays using FITC-labelled protein       28         5.19. Liquid proteolytic (alcalase) activity assays using p-nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using p-nitroanalide chromophores       29         5.21. Liquid hydrophilic and dyeability modification of fabrics       29         5.22. Liquid hydrophilic and dyeability modification of fabrics       29         5.23. Liquid amino modification of polyamide fabrics       30         5.24. Dye liquid discoloration test       31         6. Protocols relevant for COSMETIC APPLICATIONS       32         6.1. Liquid colorimetric assay for hyaluronidase activity       33         6.1.2. BSA method       33         6.1.4. DMAB method       34         6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD       34         6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD       34		5.11.1. NEFA-HR(2) kit	. 25
5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC.       27         5.15. Liquid oxidoreductase (laccase/oxidase) quantification assay       27         5.16. Liquid amylase quantification assay with chromogenic substrates       28         5.17. Liquid amylase quantification assay with polymeric substrates       28         5.18. Liquid proteolytic activity assays using FITC-labelled protein       28         5.19. Liquid proteolytic (alcalase) activity assays using <i>p</i> -nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores       29         5.21. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores       29         5.22. Liquid hydrophilic and dyeability modification of fabrics       29         5.23. Liquid anino modification of polyamide fabrics       30         5.24. Dye liquid discoloration test       31         6. Protocols relevant for COSMETIC APPLICATIONS       32         6.1. Liquid colorimetric assay for hyaluronidase activity       33         6.1.2. BSA method       33         6.1.4. DMAB method       34         6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD       34         6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD       34		5.12. Liquid polyesterase quantification assay with polylactic acid (PLA) powder	27
5.15. Liquid oxidoreductase (laccase/oxidase) quantification assay275.16. Liquid amylase quantification assay with chromogenic substrates285.17. Liquid amylase quantification assay with polymeric substrates285.18. Liquid proteolytic activity assays using FITC-labelled protein285.19. Liquid proteolytic (alcalase) activity assays using <i>p</i> -nitroanalide chromophores295.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores295.21. Liquid proteolytic activity assays using azocasein295.22. Liquid hydrophilic and dyeability modification of fabrics295.23. Liquid anino modification of polyamide fabrics305.24. Dye liquid discoloration test316. Protocols relevant for COSMETIC APPLICATIONS326.1. Liquid colorimetric assay for hyaluronidase activity326.1.2. BSA method336.1.3. BCA method346.2. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		5.13. Liquid polyesterase quantification assay using Impranil DLN-SD	27
5.16. Liquid amylase quantification assay with chromogenic substrates285.17. Liquid amylase quantification assay with polymeric substrates285.18. Liquid proteolytic activity assays using FITC-labelled protein285.19. Liquid proteolytic (alcalase) activity assays using p-nitroanalide chromophores295.20. Liquid proteolytic (subtilisin) activity assays using p-nitroanalide chromophores295.21. Liquid proteolytic cuivity assays using accasein295.22. Liquid hydrophilic and dyeability modification of fabrics295.23. Liquid amino modification of polyamide fabrics305.24. Dye liquid discoloration test316. Protocols relevant for COSMETIC APPLICATIONS326.1. Liquid colorimetric assay for hyaluronidase activity326.1.2. BSA method336.1.3. BCA method346.2. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD34		5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC	27
5.17. Liquid amylase quantification assay with polymeric substrates       28         5.18. Liquid proteolytic activity assays using FITC-labelled protein       28         5.19. Liquid proteolytic (alcalase) activity assays using p-nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using p-nitroanalide chromophores       29         5.21. Liquid proteolytic (subtilisin) activity assays using p-nitroanalide chromophores       29         5.22. Liquid proteolytic activity assays using azocasein       29         5.23. Liquid amino modification of polyamide fabrics       29         5.24. Dye liquid discoloration test       30         5.24. Dye liquid discoloration test       31         6. Protocols relevant for COSMETIC APPLICATIONS       32         6.1. Liquid colorimetric assay for hyaluronidase activity       33         6.1.2. BSA method       33         6.1.3. BCA method       34         6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD       34         6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD       34         6.4. Agar assay for the determination of hyaluronidase activity       35		5.15. Liquid oxidoreductase (laccase/oxidase) quantification assay	27
5.18. Liquid proteolytic activity assays using FITC-labelled protein       28         5.19. Liquid proteolytic (alcalase) activity assays using <i>p</i> -nitroanalide chromophores       29         5.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores       29         5.21. Liquid proteolytic activity assays using azocasein       29         5.22. Liquid hydrophilic and dyeability modification of fabrics       29         5.23. Liquid amino modification of polyamide fabrics       30         5.24. Dye liquid discoloration test       31         6. Protocols relevant for COSMETIC APPLICATIONS       32         6.1. Liquid colorimetric assay for hyaluronidase activity       33         6.1.2. BSA method       33         6.1.3. BCA method       34         6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD       34         6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD       34		5.16. Liquid amylase quantification assay with chromogenic substrates	. 28
5.19. Liquid proteolytic (alcalase) activity assays using <i>p</i> -nitroanalide chromophores295.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores295.21. Liquid proteolytic activity assays using azocasein295.22. Liquid hydrophilic and dyeability modification of fabrics295.23. Liquid amino modification of polyamide fabrics305.24. Dye liquid discoloration test316. Protocols relevant for COSMETIC APPLICATIONS326.1. Liquid colorimetric assay for hyaluronidase activity326.1.1. DNS method336.1.2. BSA method346.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		5.17. Liquid amylase quantification assay with polymeric substrates	28
5.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores295.21. Liquid proteolytic activity assays using azocasein295.22. Liquid hydrophilic and dyeability modification of fabrics295.23. Liquid amino modification of polyamide fabrics305.24. Dye liquid discoloration test316. Protocols relevant for COSMETIC APPLICATIONS326.1. Liquid colorimetric assay for hyaluronidase activity326.1.1. DNS method336.1.2. BSA method346.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		5.18. Liquid proteolytic activity assays using FITC-labelled protein	28
5.21. Liquid proteolytic activity assays using azocasein295.22. Liquid hydrophilic and dyeability modification of fabrics295.23. Liquid amino modification of polyamide fabrics305.24. Dye liquid discoloration test316. Protocols relevant for COSMETIC APPLICATIONS326.1. Liquid colorimetric assay for hyaluronidase activity326.1.1. DNS method336.1.2. BSA method336.1.3. BCA method346.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		5.19. Liquid proteolytic (alcalase) activity assays using <i>p</i> -nitroanalide chromophores	. 29
5.22. Liquid hydrophilic and dyeability modification of fabrics295.23. Liquid amino modification of polyamide fabrics305.24. Dye liquid discoloration test316. Protocols relevant for COSMETIC APPLICATIONS326.1. Liquid colorimetric assay for hyaluronidase activity326.1.1. DNS method336.1.2. BSA method336.1.3. BCA method346.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		5.20. Liquid proteolytic (subtilisin) activity assays using <i>p</i> -nitroanalide chromophores	. 29
5.23. Liquid amino modification of polyamide fabrics305.24. Dye liquid discoloration test316. Protocols relevant for COSMETIC APPLICATIONS326.1. Liquid colorimetric assay for hyaluronidase activity326.1.1. DNS method336.1.2. BSA method336.1.3. BCA method346.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		5.21. Liquid proteolytic activity assays using azocasein	29
5.24. Dye liquid discoloration test316. Protocols relevant for COSMETIC APPLICATIONS.326.1. Liquid colorimetric assay for hyaluronidase activity326.1.1. DNS method.336.1.2. BSA method.336.1.3. BCA method.346.1.4. DMAB method.346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD.346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD.346.4. Agar assay for the determination of hyaluronidase activity35		5.22. Liquid hydrophilic and dyeability modification of fabrics	. 29
6. Protocols relevant for COSMETIC APPLICATIONS.       32         6.1. Liquid colorimetric assay for hyaluronidase activity       32         6.1.1. DNS method.       33         6.1.2. BSA method       33         6.1.3. BCA method.       34         6.1.4. DMAB method.       34         6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD.       34         6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD.       34         6.4. Agar assay for the determination of hyaluronidase activity       35		5.23. Liquid amino modification of polyamide fabrics	30
6.1. Liquid colorimetric assay for hyaluronidase activity326.1.1. DNS method336.1.2. BSA method336.1.3. BCA method346.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		5.24. Dye liquid discoloration test	. 31
6.1.1. DNS method.336.1.2. BSA method336.1.3. BCA method.346.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD.346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD.346.4. Agar assay for the determination of hyaluronidase activity35	6	Protocols relevant for COSMETIC APPLICATIONS	32
6.1.2. BSA method336.1.3. BCA method346.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		6.1. Liquid colorimetric assay for hyaluronidase activity	32
6.1.3. BCA method346.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		6.1.1. DNS method	33
6.1.4. DMAB method346.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD346.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD346.4. Agar assay for the determination of hyaluronidase activity35		6.1.2. BSA method	33
<ul> <li>6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD</li></ul>		6.1.3. BCA method	34
<ul><li>6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD</li></ul>		6.1.4. DMAB method	34
6.4. Agar assay for the determination of hyaluronidase activity		6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD	34
		6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD	34
Annex - List of esters successfully tested with pH shift assay		6.4. Agar assay for the determination of hyaluronidase activity	35
	A	nnex - List of esters successfully tested with pH shift assay	36

## Abbreviations

4-AA, 4-aminoantipyrine  $\epsilon$ , Extinction coefficient ABTS, 2,2'-Azinobis(3-ethylbenzthiazoline-6-sulfonate) ACOD, Acyl-CoA oxidase ACS, Acyl-CoA synthetase AMP, adenosine monophosphate ATP, adenosine 5-triphosphate BCA, Bicinchoninic acid BES, N,N-bis(2-hydroxyethyl)-2-aminoethanesulfonic acid BHET, bis-(2-Hydroxyethyl) terephthalate C12:0, Trilaurin C14:0, Trimyristin C16:0, Tripalmitin C16:1, Tri-glyceryl palmitoleic acid C17:0, Tri-glyceryl heptadecanoic acid C18:0, Tristearate C18:1, Triolein C18:2, Trilinolein CHES, N-Cyclohexyl-2-aminoethanesulfonic acid CoA, Coenzyme A CO, Cotton DAS, 3-Amino-5-nitrosalicylic acid DMAB, p-Dimethylaminobenzaldehyde DMSO, dimethyl sulfoxide DNS, 3,5-Dinitrosalicylic acid EPPS, 4-(2-Hydroxyethyl)-1-piperazinepropanesulfonic acid FITC, Fluorescein isothiocyanate FRET, Fluorescence resonance energy transfer GlcNAc, N-Acetyl-D-glucosamine HAOS, Hyaluronic acid oligosaccharides HEPES, 4-(2-Hydroxyethyl)-1-piperazineethanesulfonic acid () HPAEC-PAD, High performance anion exchange chromatography with pulsed amperometric detection HPLC-SEC, High performance size exclusion chromatography with evaporative light scattering detector LB, Luria Bertani MEHA, 3-Methyl-N-Ethyl-N-(β-Hydroxyethyl)-Aniline MHET, Mono-(2-hydroxyethyl) terephthalate MWCO, Molecular weight cut-off NEFA, Non-esterified fatty acid PA, Polyamide fibres PCD, Polycaprolactone diol PCL, Polycaprolactone PES, Polyester fibres PET, Polyethylene terephthalate PLA, Poly(DL)-lactide or polylactic acid p-NP, p-Nitrophenol Ppi, pyrophosphoric acid RFU, Relative fluorescence units TB, Terrific broth TBS, Tris buffered saline TNBS, Trinitrobenzenesulfonic acid TPA, Terephthalic acid WR, Working reagent

5

# 1. Scope of the Deliverable

This deliverable consists in a report detailing the customized multi-substrate and multi-condition screen protocols to be applied by partners to speed up the screening of bio-resources for enzymes, and further characterization of enzymes and their engineered variants, of interest for the purposes of FuturEnzyme's objectives. On the basis of the activity level units, decisions are taken to pre-select those best matching industrial requirements to proceed with sequencing and characterisation. The protocols are available in the internal FuturEnzyme repository.

## 2. Reasons for the update

The first version of the Deliverable D3.2 was submitted in July 2021. This update is due to the fact that in these approx 18 months (from July 2021-original submission- to December 2022 -update submission), the partners have exchanged new materials for tests from the manufacturers and from commercial suppliers, and new protocols to evaluate enzymatic activity have been defined. In November 2022, the Coordinator (Manuel Ferrer) contacted the Project Officer (Colombe Warin) to explain these circumstances and ask her to re-open the submission of this deliverable (amongst others), at which she agreed.

# 3. Origin of the Deliverable

Along the already 18 months of project, one deliverable has been accomplished from which the present one nourishes. To be mentioned:

• D2.1: Manufacturers' needs and specifications: protocol (August 2021, to be updated by December 2022) In this deliverable, information about manufacturers' needs, and enzymes and products specifications (working/storage conditions and stabilities, compositions, etc.) for implementing 3 innovative, real-life, and environment-friendly products (detergents, textiles and consumer care products) are detailed. They include: Detergents: Enzymes for removing fatty oil stains Cosmetic: Enzymes for degrading hyaluronic acid to size controlled products to be integrated into cosmetics Textile: Enzymes for the removal of spinning additives and dyes

Based on these applications and needs detailed in D2.1, a series of protocols and assays have been established to help not only to identify enzymes or microorganisms that contain them, but also to assist in their detailed characterization in order to identify the best candidates. The following are the protocols and assays developed and used by the different partners during the project. Some of these protocols are part of prior knowledge, but others have been designed and adapted to the needs of the project.

# 4. Protocols for DETERGENT APPLICATIONS

As detailed in Deliverable D2.1, priority enzymes are those for removing fatty oil stains, that will include:

- True lipases (EC 3.1.1.3)
- Esterases (EC 3.1.1.1)
- Cutinases (EC 3.1.1.74) and related fatty-oil degrading hydrolases

The enzymes should be active and stable under conditions relevant to the wash cycle and to storage. Below, the key specifications are summarized:

• The enzymes should be stable for at least 2 to 3 months at 30°C in the liquid detergent formulation provided by partner HENKEL (see details in Deliverable D2.1).

• The enzymes should be effective and stable at a washing temperature between 20 and 40°C and at pH 7.0-8.5, at least during an operation time of a common wash cycle (120 min). Note: These stability and activity refer to that of the enzymes in a wash liquor mimicking the detergent-water mixture in a wash machine; this wash liquor consists in about 50 g liquid detergent per 20 L of water.

Therefore, it is a priority to adapt the screening and characterization methods to those enzymes and conditions. Below, a number of protocols are described which have been proven to successfully work in a number of buffers and conditions. As detailed in Deliverable D2.1, in general Henkel strongly recommends to concentrate on the screening methods which can be performed in a wash-liquor matrix (instead of standard buffers) as early as possible, since this affects the enzyme properties often quite strongly. It is similarly crucial to screen on real textiles as soon as possible, too, as they are more challenging than stains alone. Therefore, it is highly recommended that partners focus on, and adapt the screen methods to, (A) commercially available standard soils on textiles and (B) natural soils of interest with high consumer relevance detailed in Deliverable D2.1. Note: priority standard soil textiles recommended by Henkel, as detailed in Deliverable D2.1, have been ordered by CSIC by July 2021 and transferred to all partners (see Deliverable D2.1) involved in enzyme screening and characterization.

The following decision-taken strategy, once activity towards ester-like substrates is confirmed, has been undertaken.

#### Decision taken screen 1:

• Enzyme retaining some activity 5-10 min in wash liquor Note: wash liquor is prepared by adding 50 g liquid detergent provided by partner Henkel per 20 L of water.

#### Decision taken screen 2:

• Active towards commercially available standard soils/stains on textiles and natural soils. They include, at least, cuff and collar, natural skin fat, butterfat, olive oil, frying fat, lard and tomato beef sauce, beef lard, pigment, oil, pigment, sebum, mayonnaise with carbon black, butterfat with colorant, lipstick, pink and make up.

Note: the following standard soils/stains on textiles and natural soils were selected by partner Henkel, which were requested to Center For Testmaterials, CFT: Pigment with oil on polyester/cotton (PC-09), Mayonnaise on cotton (C-S-05S), Lipstick, pink on polyester/cotton (P-S-16), Fluid make-up on cotton (C-S-17), High discriminative sebum BEY on polyester/cotton (PC-S-132), Beef fat on cotton (C-S-61) and Butterfat on cotton (C-S-10).

C-S-	61 Beef fat, cold	ured wit	h Suda	n red	<b>*</b> L		
	Very suitable to me Coloured with the I Not aged at elevate	east dangerou	us type of				
O de la constante	: Substrate:	Width:	Yu:	Order code:	Substrate:	Width:	
Order code				Control Contract of Contract Statistics (Section 201	Jubblidte.	width.	Yu:
C-S-61	Cotton (CN-11)	45 cm	89,1	PA-S-61	PolyAcryl (PAN-01)	45 cm	Yu: 89,3
		ALC: NOT THE REAL PROPERTY OF	89,1 94,5				Net state
C-S-61	Cotton (CN-11)	45 cm		PA-S-61	PolyAcryl (PAN-01)	45 cm	89,3

# C-S-05S

#### Mayonnaise with carbon black



For measuring the removal of high amounts of fatty matter. Additional carbon black is applied to enable assessment by reflectometry. Represents a difficult to remove composite soil containing lipids and proteins. Not aged at elevated temperature.

💧 🍁 c

G @ 🌡 -

45 cm 87,4

Order code:	Substrate:	Width:	Yu:	Order code:	Substrate:	Width:	Yu:
C-S-05S	Cotton (CN-17)	90 cm	87,8	PA-S-05S	PolyAcryl (PAN-01)	45 cm	89,3
KC-S-05S	Knitted Cotton (CN-42)	35 cm	94,5	S-S-05S	Silk (T-601)	45 cm	83,9
PC-S-05S	Polyester/Cotton (PCN-01)	90 cm	87,8	W-S-05S	Wool (T-541)	45 cm	67,2
P-S-05S	Polyester (W-30 A)	90 cm	88,4	N-S-05S	Nylon (T-365)	45 cm	87,4

#### C-09 Pigment with oil (below 60° C)



Measures general detergency. Also reacts to builder concentration, water hardness and product concentration.

Aged at elevated temperature.

Order code:	Substrate:	Width:	Yu:	Order code:	Substrate:	Width:	Yu:
C-09	Cotton (CN-11)	90 cm	89,1	PA-09	PolyAcryl (PAN-01)	45 cm	89,3
KC-09	Knitted Cotton (CN-42)	35 cm	94,5	S-09	Silk (T-601)	45 cm	83,9
PC-09	Polyester/Cotton (PCN-01)	90 cm	87,8	W-09	Wool (T-541)	45 cm	67,2
P-09	Polyester (PN-01)	90 cm	86,5	N-09	Nylon (T-365)	45 cm	87,4

C-S-1	<b>O</b> Butterfat with	c <mark>olou</mark> ra	nt		*-		
6	Dehydrated butter co of active systems and Not aged at elevated	lipase enzy	mes.	ourant. Suitable to	o evaluate performance		
Order code:	Substrate:	Width:	Yu:	Order code:	Substrate:	Width:	Yu:
C-S-10	Cotton (CN-11)	45 cm	89,1	PA-S-10	PolyAcryl (PAN-01)	45 cm	89,3
C-S-10 KC-S-10	Cotton (CN-11) Knitted Cotton (CN-42)	45 cm 35 cm	89,1 94,5	PA-S-10 S-S-10	PolyAcryl (PAN-01) Silk (T-601)	45 cm 45 cm	89,3 83,9

N-S-10

Nylon (T-365)

💧 @ :

#### C-S-16 Lipstick, pink

Polyester (PN-01)

P-S-10

For studying removal of a very realistic and common stain. Based on fatty matter and pigments. Might give redeposition problems due to high concentration of the lipstick. N

45 cm 86,5

ot aged at el	evated te	emperature.
---------------	-----------	-------------

Order code:	Substrate:	Width:	Yu:	Order code:	Substrate:	Width:	Yu:
C-S-16	Cotton (CN-11)	45 cm	89,1	PA-S-16	PolyAcryl (PAN-01)	45 cm	89,3
KC-S-16	Knitted Cotton (CN-42)	35 cm	94,5	S-S-16	Silk (T-601)	45 cm	83,9
PC-S-16	Polyester/Cotton (PCN-01)	45 cm	87,8	W-S-16	Wool (T-541)	45 cm	67,2
P-S-16	Polyester (PN-01)	45 cm	86,5	N-S-16	Nylon (T-365)	45 cm	87,4

C-S-1	L7 Fluid make-up	Ę					
0							
	Not aged at elevated						
Order code:		Width:	Yu:	Order code:	Substrate:	Width:	Yu:
C-S-17	Cotton (CN-17)	90 cm	89,1	PA-S-17	PolyAcryl (PAN-01)	45 cm	89,3
KC-S-17	Knitted Cotton (CN-42)	35 cm	94,5	S-S-17	Silk (T-601)	45 cm	83,9
PC-S-17	Polyester/Cotton (PCN-01)	90 cm	87,8	W-S-17	Wool (T-541)	45 cm	67,2

## C-S-132 High discriminative sebum BEY with pigment



Most commonly used synthetic skin fat. Improved version of C-S-32. To simulate the removal of human skin fat, using active systems and lipase systems.

00

Not aged at elevated temperature.

Order code:	Substrate:	Width:	Yu:	Order code:	Substrate:	Width:	Yu:
C-S-132	Cotton (T-400)	45 cm	88,9	PA-S-132	PolyAcryl (PAN-01)	45 cm	89,3
KC-S-132	Knitted Cotton (CN-42)	35 cm	94,5	S-S-132	Silk (T-601)	45 cm	83,9
PC-S-132	Polyester/Cotton (PCN-01)	45 cm	87,8	W-S-132	Wool (T-541)	45 cm	67,2
P-S-132	Polyester (PN-01)	45 cm	86,5	N-S-132	Nylon (T-365)	45 cm	87,4

#### Decision taken screen 3:

- Enzyme retaining activity 120 min in wash liquor
- Enzyme stable in liquid detergent for few days
- Active against standard soils/stains on textiles.

#### **Final requirements:**

- Stable 2-3 months at 30°C in the liquid detergent formulation
- Effective at conditions mimicking the wash cycle using wash liquor: 20-40°C, pH 7.0-8.5, 120 min. As a baseline or reference, a wash liquor made of "HENKEL® Liquid Laundry Detergent\_A including enzymes" is used, so that the values obtained are compared with those obtained when the FuturEnzyme's enzymes were added to a wash liquor made of "HENKEL® Liquid Laundry Detergent\_A without enzymes". Only enzymes, lipases, that behave similarly or better than the "HENKEL® Liquid Laundry Detergent\_A including enzymes" are selected.

Below, the detergents provided by partner Henkel, and that were used for activity tests when appropriated, are detailed:

#### Detergents with and without enzymes

Sent to Bangor, CSIC, UDUS, UHAM, IST-ID, CNR in September 2021

- Density: 1,0855 1,0955 g/cm3
- pH: 7,8 8,2
- Viscosity: 210 310 mPa.s

Sample name	Batch No.	Dosage for wash liquor (16.7 °dH water)	Ingredients	Gap [%] to be filled (last w water)
Detergent_A	WLHUL21 <b>61400</b>	3.1 g/L	All ingredients	-
			included	
Detergent_A_w/o	WLHUL21 <b>94600</b>	3.1 g/L	Without	2.5 %
			enzymes	

Hazardous substances according to CLP (EC) No 1272/2008:

Hazardous substances CAS-No	EINECS	REACH-Reg No.	Content	Classification
Benzenesulfonic acid, mono-C10-13-alkyl derivs., compds. with ethanolamine 85480-55-3	287-335-8	*	>= 20- < 40 %	Acute toxicity 4 H302 Skin irritation 2 H315 Serious eye damage 1 H318 Chronic hazards to the aquatic environment 3 H412
Alcohols, C12-18, ethoxylated 68213-23-0			>= 10- < 20 %	Acute toxicity 4 H302 Serious eye damage 1 H318 Chronic hazards to the aquatic environment 3 H412
Alcohols, C12-14, ethoxylated, sulfates, sodium salts 68891-38-3	500-234-8	01-2119488639-16	>= 5- < 10 %	Skin irritation 2; Dermal H315 Serious eye damage 1 H318 Chronic hazards to the aquatic environment 3 H412
Tetrasodium (1- hydroxyethylidene)bisph osphonate 3794-83-0	223-267-7	*	>= 1- < 5 %	Acute toxicity 4 H302 Serious eye irritation 2 H319
Sodium metaborate, anhydrous 7775-19-1	231-891-6	*	>= 1- < 5,9 %	Toxic to reproduction 2 H361d Serious eye irritation 2 H319
Protease 9014-01-1	232-752-2	01-2119480434-38	>= 0,1- < 1 %	Acute hazards to the aquatic environment 1 H400 Chronic hazards to the aquatic environment 2 H411 Acute toxicity 4 H302 Specific target organ toxicity - single exposure 3 H335 Skin irritation 2 H315 Serious eye damage 1 H318 Respiratory sensitizer 1 H334

\*exempted according to REACH article 2(7) and Annex V. Each starting material of the ionic mixture is registered, as required.

## 4.1. pH shift liquid protocols for quantifying esterase-lipase activity

The protocols are recommended for short-middle chain length carboxylesterases, and have been successfully applied for a variety of chemically and structurally diverse soluble and insoluble esters, including short-medium-large triglycerides (see Annex). The method allows using insoluble triglycerides.

## 4.1.1. Using Phenol Red (pH 8.0) as a pH indicator

Below the assay is described for 384-well plates, but can be adapted to 96-well plates or any other format (e.g. 1 mL cuvette).

- Prepare a concentrated ester stock solution<sup>1</sup> by dissolving each ester at a concentration of 100 mM in acetonitrile, dimethyl sulfoxide (DMSO), acetone or isopropanol, depending on its solubility (*stock solutions can be prepared and maintained at -20°C until use*).
- Fill a 384-well plate with 20 μL of 5 mM 4-(2-hydroxyethyl)-1-piperazinepropanesulfonic acid (EPPS) buffer pH 8.0<sup>2</sup>, using a QFill3 microplate filler.
- Add 2 µL of each ester stock solution to each well using a PRIMADIAG liquid-handling robot.
- After the esters are added, fill the 384-well plate with 20 μL of 5 mM EPPS buffer pH 8.0, containing 0.912 mM Phenol Red (used as a pH indicator: from red to yellow color when pH is lowered) using a QFill3 microplate filler (*the final ester concentration in each well is 4.5 mM, and the final concentration of Phenol Red is 0.45 mM*).
- Add immediately 2 μL of enzyme solution, cell extract or cell suspension<sup>3</sup> to each well using an Eppendorf Repeater M4 pipette.
- Follow the progress of the reaction at the desired temperature by monitoring the absorbance at 550 nm<sup>4</sup>, in continuous mode, in a microplate reader (*reaction can be followed for up to 24 h, but we recommend continuous reading for up to 1-8 hours, depending on the enzyme activity*).
- Activity (unit/mg protein) is calculated by determining the absorbance per minute from the slopes generated, and by applying the following formula:

 $Rate\left(\frac{\mu mol}{\min mg \ prot./cells}\right) = \frac{\frac{\Delta Abs}{\min}}{8450 \ M - 1 \text{cm} - 1} * \frac{1}{0.4 \ cm} * \frac{10^6 \ \mu M}{1 \ M} * 0.000044 \ L * \frac{1}{mg \ prot./cells}$ 

Extinction coefficient ( $\epsilon$ ) phenol red at 550 nm and pH 8.0: 8450  $M^{\text{-1}}\text{cm}^{\text{-1}}$  Total assay volume: 44  $\mu\text{L}$ 

<sup>1</sup>This method has been proven efficient for a wide variety of chemically and structurally diverse esters, including long chain triglycerides and poly-aromatic esters.

<sup>2</sup>This method can be adapted to the use of wash liquor supplied by Henkel. Briefly, the 5 mM EPPS buffer pH 8.0, is substituted by one in which 0.5 g liquid detergent is added to 100 ml of 5 mM EPPS buffer pH 8.0. Alternatively, when using large triglycerides such as those based on C12:0 (trilaurin), C14:0 (trimyristin), C16:0 (tripalmitin), C16:1 (triolein), C17:0, C18:0, C18:1, and C18:2 (trilinolein), the 5 mM EPPS buffer pH 8.0, can be substituted by one containing 0.2% (w/w) arabic gum and 0.4% (w/v) sodium deoxycholate.

<sup>3</sup>Cell suspension can be prepared following optimal cultivation conditions. In case of *Escherichia coli* expressing enzymes of interest the following protocol is recommended. We do the cultivation in 96 well deep-well-plates, starting with 300  $\mu$ L Terrific Broth (TB) medium with the required antibiotic. Wells get inoculated with a loop of the cryoculture or an single colony from an agar plate and grown over night at 37°C, 200 rpm, and sealed with a breathable membrane. The next day, 3  $\mu$ L of each preculture are transferred to fresh 300  $\mu$ L TB + antibiotic and grown for 3 h, 37°C, 200 rpm. Then cultures get induced with 1 mM IPTG (or other appropriate inducer) and grown over night at 30°C, 200 rpm. Next, the cultures are spun down in the plate, the media is removed and plates are covered with a fresh breathable membrane and frozen for at least 6 h at -30°C. Finally the plates get lyophilized overnight. For assays, the liophilized cells are resuspended in 0.6 ml of 5 mM EPPS buffer pH 7.0 and 2  $\mu$ l are used for activity tests

<sup>4</sup>The acid produced after ester bond cleavage by the hydrolytic enzyme induces a color change in the pH indicator that is measured spectrophotometrically at 550 nm. Absence of activity defined as at least a 2-fold background signal.

## 4.1.2. Using *p*-nitrophenol (pH 7.0) as a pH indicator

- Prepare the reaction mixture in 96-well microplates adding 150 μl of 2 mM N,N-bis(2-hydroxyethyl)-2aminoethanesulfonic acid (BES) buffer pH 7.2 containing 0.45 mM *p*-nitrophenol
- Add 2 mM ester substrate (soluble aliphatic and aromatic esters, available from Merck/Sigma)<sup>1</sup>
- Add 0.1-0.5  $\mu g$  of purified protein or about 2  $\mu L$  enzyme solution, cell extract or cell suspension to each well.
- Incubate the plates for 10 min at the desired temperature (e.g., 25-37°C).
- Monitore spectrophotometrically the hydrolytic activity by following absorbance of *p*-nitrophenol at 404 nm (ε = 17.3 mM<sup>-1</sup> cm<sup>-1</sup>) (Janes L.E. et al., 1998, Chem. Eur. J., 4: 2324-2331).
- Activity (unit/mg protein) is calculated by determining the absorbance per minute from the slopes generated, and by applying the following formula:

$$Rate \left(\frac{\mu mol}{\min mg \ prot./cells}\right) = \frac{\frac{\Delta Abs}{\min}}{17300 \ M - 1 \text{cm} - 1} * \frac{1}{0.4 \ cm} * \frac{10^{6} \ \mu M}{1 \ M} * 0.00015 \ L * \frac{1}{mg \ prot./cells}$$

- Extinction coefficient (ε) *p*-nitrophenol at 404 nm and pH 7.2: 17300 M<sup>-1</sup>cm<sup>-1</sup>
- Total assay volume: 150 μL

<sup>1</sup>A 100 mM stock solution of the esters can be prepared in acetonitrile, dimethyl sulfoxide (DMSO), acetone or isopropanol, depending on their solubility.

#### 4.1.3. Using nitrazine yellow as a pH indicator

The method is adapted to handle insoluble substrates (semi-quantitative) and allows high concentrations of co-solvents.

Harvest the expression cultures by centrifugation, discard the supernatants, and resuspend the cells in 0.1 mL cell lysis solution containing polymyxin B (10 mM potassium phosphate buffer pH 7.2, 0.1 mg/mL Polymyxin B).

- Incubate for 1h at 37°C.
- For solvent tolerance assessment: mix in a 96 well plate 0.1 mL of the esterase-containing cell extracts with 0.1 mL of the respective solvent/water mix in a microplate to reach final solvent concentrations of 0%, 30%, and 50% (vol/vol) and incubate for 2 h at 30°C. During this pre-incubation, the microplate lid should be sealed with organic-solvent-stable tape to prevent evaporation.
- Combine in a 96 well plate 10 μL of the sample (cell extract or organic solvent pre-incubation mix) with 180 μL nitrazine yellow-containing assay buffer (5 mM potassium phosphate buffer pH 7.2, 20 μg/mL nitrazine yellow, and 0%, 30%, or 50% (vol/vol) of the respective organic solvent) and 10 μL of substrate solution (e.g. 200 mM tributyrin in acetonitrile) or 10 μL acetonitrile for the control. In the case of a color shift after the addition of the organic solvent and before substrate supplementation, the pH is titrated to neutral (blue color) with potassium hydroxide solution.
- Incubate the reaction mixture for 18 h at 30°C.
- Determine the activity by measuring the absorption at 450 and 600 nm. The quotient of the absorption values determined at both wavelengths is used to measure the pH shift. Correct each value by subtraction of the control which did not contain substrate before calculation of mean values and standard deviations.

To reduce false positives, values in the range of the standard deviation of the empty vector control should be considered as not active.

## 4.1.4. Using fluorescein as pH indicator

The method is adapted to handle insoluble substrates (semi-quantitative) and allows usage of solid substrates.

- Prepare 2 mM potassium phosphate (KPi) buffer pH 7.0 or Tris-buffer pH 8.0 (this buffer can be adapted to specific requirement).
- Prepare working reagent; buffer + 5.0 μM Fluorescein (+ substrate dispersion, if applicable. For Impranil DLN, we use typically 5 μL/mL).
- Add 0.19 mL working reagent (and substrate) per well of a 96-well plate.
- Add 10  $\mu$ L of enzyme solution to the wells (and 10  $\mu$ L of the buffer as control).
- Follow the activity (= drop of fluorescence) at Excitation: 488 nm/Emission: 525 nm at the temperature of interest.
- If the measurement is performed longer than 15 min, add a cover to the plate to prevent evaporation.

## 4.2. Liquid protocol for quantifying esterase-lipase activity with chromogenic esters

The protocols are recommended for short-middle chain length carboxylesterases.

## 4.2.1. Using short-medium $\alpha$ - or $\beta$ -naphthyl and *p*-nitrophenol esters

- Prepare the reaction mixture (200 μL) containing 50 mM Tris-HCl buffer pH 8.0 or N-cyclohexyl-2aminoethanesulfonic acid (CHES) buffer pH 9.0-10.0, 1 mM substrate (naphthyl) acetate, butyrate, laurate, etc., and 4-nitrophenyl esters such as *p*NP-butyrate (C4), caprylate (C8), laurate (C12), palmitate (C16), and stearate (C18)], and 0.01-1.0 μg of purified enzyme or appropriated amount of cell extracts or cell suspensions.
- Incubate the reaction for 10 min at selected temperatures (from 15 to 50°C).
- Monitore spectrophotometrically the activity by following absorbance at 310 nm (for  $\alpha$  or  $\beta$ -naphthyl esters of fatty acids substrates) or 410 nm (for *p*-nitrophenyl (*p*NP) substrates).

## 4.2.2. Using short-large *p*-nitrophenol esters

- Prepare the substrate stock with 20 mM 4-nitrophenyl ester [*p*NP-butyrate (C4), caprylate (C8), laurate (C12), palmitate (C16), and stearate (C18)]) in acetonitrile.
- Prepare working reagent of 100 mM potassium phosphate buffer pH 7.2, 1 mM 4-nitrophenyl ester, 5% (vol/vol) acetonitrile from 20 mM substrate in acetonitrile stocks.
- Pipette 10 μL of enzyme solution (vol. sample) into the well of a 96 well plate.
- Add 190  $\mu$ L of working reagent to each well (total vol. in microtiter plate (MTP) = 200  $\mu$ L).
- Measure the increase of absorption at 410 nm at 30°C for 10 min.
- Calculate activity with

 $\frac{slope \ at \ 410 \ nm \ [min^{-1}] \ * \ total \ vol. \ in \ MTP \ [ml] \ * \ dilution \ factor}{path \ length \ [cm] \ * \ \varepsilon \ [mM^{-1} \ * \ cm^{-1}] \ * \ vol. \ sample \ [ml]}$ 

and  $\epsilon$  =10 mM  $^{\text{-1}}\text{cm}^{\text{-1}}$ 

## 4.2.3. Using medium-large *p*-nitrophenol esters

- Prepare working reagent of 25 mM Tri-HCl buffer, pH 7.0, containing 0.1% (w/w) arabic gum and 0.2% (w/v) sodium deoxycholate at 37°C. Add 4-nitrophenyl ester [*p*NP-butyrate (C4), caprylate (C8), laurate (C12), palmitate (C16), and stearate (C18)] to a final concentration of 1-8 mM.
- Pipette 190 μL of the working reagent into the well of a 96 well plate.
- Pipette 10 µL of enzyme solution (vol. sample) into the well of a 96 well plate.
- Measure the increase of absorption at 410 nm at 30°C for 10 min, and calculate the activity as above.

## 4.3. Liquid protocol for quantifying esterase-lipase activity with non-chromogenic esters

#### 4.3.1. Using with fatty acid cholesterol esters

The protocols are recommended for short-middle-large chain length carboxylesterases.

- Prepare working reagent of 25 mM Tri-HCl buffer pH 7.0, containing 0.1% (w/w) arabic gum and 0.2% (w/v) sodium deoxycholate at 37°C. Then, add cholesteryl oleate<sup>1</sup> to a final concentration of 1 mM.
- Pipette 190  $\mu L$  of the working reagent into the well of a 96 well plate.
- Pipette 10 µL of enzyme solution (vol. sample) into the well of a 96 well plate.
- Keep reaction for 20 min and estimate the cholesterol released by a commercial enzymatic method (Boehringer).
- One unit of activity was defined as the amount of enzyme hydrolyzing 1  $\mu mol$  of substrate per minute under the above conditions.

<sup>1</sup>This method can be used for testing cholesteryl butyrate (C4:0), cholesteryl palmitate (C16:0), cholesteryl stearate (C18:0), cholesteryl oleate (C18:1) and cholesteryl linoleate (C18:2).

## 4.3.2. Using insoluble substrates (waxes and lipids) and pH-stat

The hydrolysis of tributyrin, triolein, cholesterol esters, and other waxes and lipids can be assayed titrimetrically at pH 7.0 and 25°C in a pH-stat (Mettler, model DL50) using 0.1 M NaOH as titrant.

- Prepare a reaction mixture (20 ml) containing the substrate (50 mM) in the presence of 0.15 M NaCl and 5% (v/v) Genapol X-100. Introduce this mixture into a pH-stat.
- Pipette 10-100 µL of enzyme solution (vol. sample).
- Follow hydrolysis titrimetrically at pH 7.0 and 25°C in a pH-stat using 0.1 M NaOH as titrant.
- Activity (unit/mg protein) is calculated by determining the slope of the line.

This protocol could be adapted to conditions mimicking those in a washing machine and can be adapted to any kind of insoluble substrate (wax, lipid, etc.). Instead of using commercial sufactants or dispersants such as arabic gum or Genapol X-100, the insoluble lipids are resuspended in a wash liquour mimicking the detergent-water mixture in a washing machine.

- Prepare a wash liquor containing 0.5 g liquid detergent (Henkel) per 200 mL of water.
- Prepare the reaction mixture by adding to the wash liquor the raw substrate (wax, lipid) to a final concentration of approx. 0.1-1.0 % (w/v).
- Take 20 mL of the wash liquid and incorporate it into the vial of the pH-stat.
- Pipette 10-100 μL of enzyme solution (vol. sample).
- Follow hydrolysis titrimetrically at pH 7.0 and 25°C in a pH-stat using 0.1 M NaOH as titrant.
- Activity (unit/mg protein) is calculated by determining the slope of the line.

## 4.4. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits

## 4.4.1. NEFA-HR(2) kit

The NEFA-HR(2) kit (Waco, 517982 - Reagent 1 Set kit NEFA-HR(2) Assay, 517984 - Reagent 2 Set, kit NEFA-HR(2) Assay, 517954 - Standard, kit NEFA-HR(2) Assay), also referred to as ACS-ACOD method, is used for the quantitative determination of Non-Esterified Fatty Acids (NEFA). Non-esterified fatty acid (NEFA) in sample is converted to Acyl-CoA, adenosine monophosphate (AMP) and pyrophosphoric acid (PPi) by the action of Acyl-CoA synthetase (ACS), under coexistence with coenzyme A (CoA) and adenosine 5-triphosphate disodium salt (ATP). Obtained Acyl-CoA is oxidized and yields 2,3-trans-Enoyl-CoA and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) by the action of Acyl-CoA oxidase (ACOD). In the presence of a peroxidase, the H<sub>2</sub>O<sub>2</sub> formed yields a blue-purple pigment by quantitative oxidation condensation with 3-Methyl-N-Ethyl-N-( $\beta$ -Hydroxyethyl)-Aniline (MEHA) and 4-aminoantipyrine (4-AA). Non-esterified fatty acids (NEFA) concentration is obtained by measuring absorbance of the blue-purple color.


#### Reagent preparation:

- R1: Prepare R1 by mixing one bottle of Color A and Solvent A. After preparing the R1, store at 2 10°C and use within 1 month.
- R2: Prepare R2 by mixing one bottle of Color B and Solvent B. After preparing the R2, store at 2 10°C and use within 1 month.



Protocol for stained clothes (adapted from the kit specifications):

- Incubate the enzyme (at different concentrations) with a small piece (4 mg) of a stained cloth (e.g. purchased from Center For Testmaterials, CFT), previoulsy cutted using a circle hole puncher (5 mm diameter) in 100 μL of 40 mM of the appropriated buffer (e.g. pH 7.0 HEPES buffer) or washing liquor (e.g. appropriated dilution from the detergent provided by Henkel) for 24h, at 30°C and 950 rpm. Two "small wash tests" were performed: one with with HENKEL<sup>®</sup> Liquid Laundry Detergent without enzymes (2.5 g/L water), and one with HENKEL<sup>®</sup> Liquid Laundry Detergent\_A including enzymes (3.1 g/L water).
- The reaction outcome is tested using the NEFA-Kit, following specifications:
  - Transfer 10 μL of the reaction solution + 100 μL of NEFA solution 1 to a 96 microplate well. Incubate 6 min at 37°C.
  - Add 50 μL of NEFA solution 2. Incubate 6 min at 30°C.
  - Measure sample's absorbance at 550 nm using a microplate spectrometer (e.g. Synergy HT Multi Mode Microplate Reader, Agilent, Madrid, Spain).



#### A summary of the protocol is detailed, in a schematic manner, below:

#### 4.5. Agar plate for screening esterase-lipase with $\alpha$ -naphthyl ester

This protocol is based on the use of the substrate  $\alpha$ -naphthyl acetate or laurate that is enzymatically hydrolyzed by esterases/lipases, liberating a free naphthol product. This then couples with a diazonium compound (Fast Blue RR solution), forming brown colored deposits; the positive clones will appear due to the formation of a dark brown precipitate.

- Prepare 40 mM 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) buffer pH 7.0 from 1 M stock (*Note: any other buffer can be used*). Add agar to a final concentration 0.4% and heat it in the microwave (defreeze position) to dissolve the entire agar. Cool down to ~50°C.
- Prepare 40-80 mg/mL Fast Blue RR (Sigma-Aldrich/Merck) solution in DMSO (FBRR solution).
- Prepare 20 mg/mL α-naphtyl acetate or laurate (Sigma-Aldrich/Merck) solution in acetone (*substrate solution*).
- Add 320 μL of *FBRR solution* and 320 μL of *SUBSTRATE solution* to a Falcon tube containing 20 mL of HEPES/agar previously tempered, and mix by inverting the tube.
- Overlay the plates containing individual clones or microbial colonies with 10-15 mL of the mix.
- Positive clones will appear as dark brown colonies after 30 sec to 60 min and can be picked (no longer than 5 min, if colonies do not change to dark brown color.

## 4.6. Agar plate tributyrin screening esterase-lipase assay

The protocol is recommended for short-chain length carboxylesterases.

- Prepare a 50% (v/v) tributyrin (Applichem, Darmstadt, Germany) emulsion in sterile distilled water and add 50 g/L arabic gum (Carl Roth) (Jaeger and Kovacic, 2014). Arabic gum powder is used as an emulsifying agent for the triglyceride.
- Homogenize the mixture for at least 1 min to yield a stable emulsion, e.g. using an Ultra Turrax<sup>1</sup>.
- Add 30 mL of tributyrin emulsion per 1 L of molten Luria Bertani (LB) agar and mix thoroughly, e.g. using an Ultra Turrax.
- Pour 25 mL medium portions into appropriate Petri dishes and allow the agar to solidify for at least 15 min.
- Plate bacterial clones and incubate at optimal growth temperature for the specific organism for at least 16 h.

<sup>1</sup>Add the arabic gum powder to the respective volume of water. Filling water into a tube or a bottle with a layer of the powder at the bottom should be avoided because it will result in a hard-to-dissolve clot of gum.

## 4.7. Agar plate coconut oil screening lipase assay

The protocol is recommended for middle-chain length carboxylesterases.

- Melt coconut oil (Biozentrale Naturprodukte, Wittibreut Ulbering, Germany) by incubation at 30-37°C. Pre-heat sterile distilled water to 60°C<sup>1</sup>.
- Prepare a 50% (v/v) coconut oil (e.g., Sigma-Aldrich/Merck, ref. C1758-100G) emulsion in the pre-heated water containing 50 g/L arabic gum (Carl Roth) and 0.35 g/L rhodamine B (Sigma-Aldrich/Merck, Darmstadt, Germany). Homogenize the mixture for at least 1 min to yield a stable emulsion<sup>2,3</sup>.
- Add 20 mL of coconut oil emulsion per 1 L of molten LB agar and mix thoroughly, e.g. using an Ultra Turrax.
- Pour 25 mL medium portions into appropriate Petri dishes and allow the agar to solidify for at least 15 min.
- Plate bacterial clones and incubate at optimal growth temperature for the specific organism for at least 16 h to develop fluorescent halos/fluorescent colonies.

<sup>1</sup>Heating the water in advance should avoid a drop of temperature below 30°C during the preparation of the emulsion in the next step and therefore prevent a partial hardening of the coconut oil which will hamper successful emulsification <sup>2</sup>Add the arabic gum powder to the respective volume of water. Filling water into a tube or a bottle with a layer of the powder at the bottom should be avoided because it will result in a hard-to-dissolve clot of gum.

<sup>3</sup>This method can be adapted to the use of wash liquor supplied by Henkel. Briefly, the 50% (v/v) cocoa butter triglycerides emulsion containing 50 g/L arabic gum (Carl Roth) and 0.35 g/L rhodamine B (Sigma-Aldrich/Merck, Darmstadt, Germany), could be substituted by one 50% (v/v) cocoa butter triglycerides emulsion containing 0.25 g liquid detergent supplied by Henkel per 100 ml water.

## 4.8. Agar plate cocoa butter screening lipase assay

The protocol is recommended for middle-large chain length carboxylesterases.

- Melt cocoa butter triglycerides (Sigma-Aldrich/Merck, ref. IRMM801-5G) by incubation at 30-37°C. Preheat sterile distilled water to 60°C<sup>1</sup>.
- Prepare a 50% (v/v) cocoa butter triglycerides emulsion in the pre-heated water containing 50 g/L arabic gum (Carl Roth) and 0.35 g/L rhodamine B (Sigma-Aldrich/Merck, Darmstadt, Germany). Homogenize the mixture for at least 1 min to yield a stable emulsion<sup>2,3</sup>.

- Add 20 mL of cocoa butter triglycerides emulsion per 1 L of molten LB agar and mix thoroughly, e.g. using an Ultra Turrax.
- Pour 25 mL medium portions into appropriate Petri dishes and allow the agar to solidify for at least 15 min.
- Plate bacterial clones and incubate at optimal growth temperature for the specific organism for at least 16 h to develop fluorescent halos/fluorescent colonies.

<sup>1</sup>Heating the water in advance should avoid a drop of temperature below 30°C during the preparation of the emulsion in the next step and therefore prevent a partial hardening of the coconut oil which will hamper successful emulsification <sup>2</sup>Add the arabic gum powder to the respective volume of water. Filling water into a tube or a bottle with a layer of the powder at the bottom should be avoided because it will result in a hard-to-dissolve clot of gum.

<sup>3</sup>This method can be adapted to the use of wash liquor supplied by Henkel. Briefly, the 50% (v/v) cocoa butter triglycerides emulsion containing 50 g/L arabic gum (Carl Roth) and 0.35 g/L rhodamine B (Sigma-Aldrich/Merck, Darmstadt, Germany), could be substituted by one 50% (v/v) cocoa butter triglycerides emulsion containing 0.25 g liquid detergent supplied by Henkel per 100 ml water.

## 4.9. Agar plate olive oil lipase screening assay

The protocol is recommended for long-chain length carboxylesterases.

- Prepare a 50% (v/v) olive oil<sup>1</sup> emulsion in water containing 50 g/L arabic gum (Carl Roth) and 0.35 g/L rhodamine B (Sigma-Aldrich/ Merck, Darmstadt, Germany).
- Homogenize the mixture for at least 1 min to yield a stable emulsion.
- Add 20 mL of olive oil emulsion per 1 L of molten LB agar and mix thoroughly, e.g. using an Ultra Turrax.
- Pour 25 mL medium portions into appropriate Petri dishes and allow the agar to solidify for at least 15 min.
- Plate bacterial clones or specific organism and incubate at optimal growth temperature for at least 16 h to develop fluorescent halos/fluorescent colonies.

<sup>1</sup>This method can be use also for oils such as sunflower oil, coconut oil, cocoa butter, etc.

## 4.10. Agar plate palm oil screening lipase assay

The protocol is recommended for long-chain length carboxylesterases.

- Prepare a 50% (v/v) palm oil<sup>1</sup> in water containing 50 g/L arabic gum (Carl Roth).
- Homogenize the mixture for at least 1 min to yield a stable emulsion.
- Add 20 mL of palm oil emulsion per 1 L of molten LB agar and mix thoroughly, e.g. using an Ultra Turrax.
- Pour 25 mL medium portions into appropriate Petri dishes and allow the agar to solidify for at least 15 min.
- Plate bacterial clones or specific organism and incubate at optimal growth temperature for at least 16 h, but preferably for 24-72 h.
- After cultivation, plates are overlaid with a 25 ml of a solution prepared as follows: 0.4% w/v of agar is added to a 5 mM EPPS buffer pH 8.0 containing 0.45 mM Phenol Red (used as a pH indicator) and 0.4% w/v of agar (Note: heat it in the microwave to dissolve the entire agar). After cooling down to ~40-50°C, the plate is overlaid with this solution and yellow halo zones seen are considered as positive for lipase production.

<sup>1</sup>This method can be used with any wax or lipid, including long chain triglycerides.

## 4.11. Agar plate egg yolk phospholipase assay

The protocol uses the phospholipids from egg yolk to detect phospholipase activity. Hydrolysis of phospholipids drives to the liberation of fatty acids. Calcium forms a complex with the free fatty acids resulting in the formation of a precipitation halo around the colonies. Colonies are grown in this medium.

- Add an egg yolk (~ 8% v/v) to each 500 mL of appropriate agar medium (e.g. LB agar) cooled to less than 40°C after autoclaving. Then, add appropriate antibiotics and supplements such as inductors.
- Mix vigorously using a vortex before plating into Petri dishes (mix vigorously the bottle containing the agar medium before plating every 4-5 plates or as many times as necessary).
- Replicate the clones or microbial cells into egg yolk agar plates.
- Incubate the plates for 16 hours at 37°C.
- Positive clones will appear as colonies surrounded by a halo.

## 4.12. Liquid assay for investigating effect of laundry detergents

Henkel will provide a detergent without enzymes.

- The detergent will be diluted in double distilled water to a final concentration of 2.5 mg/mL to simulate washing conditions. The enzyme is added to the reaction mix and incubated at 20-40°C for 30-120 min. Any of the method described above can be used.
- For stability tests, aliquots are taken at different time intervals and the residual activity is determined at 20°C; the enzyme will be added to different amount in the liquid detergent and incubated for months at 20-40°C. At different time frames, aliquots are taken and the activity evaluated using any of the methods described above.

## 4.13. Protocol for preparing cell suspensions for activity assays

In some cases, the presence of lipase activity or the analysis of substrate specificity can be evaluated by using cells containing presumptive lipase activity, not purified protein. The cell suspensions can be prepared following optimal cultivation conditions. In case of *Escherichia coli* expressing enzymes of interest the following protocol is recommended.

- Cultivate the cells in 96 well deep-well-plates, starting with 300 μL Terrific Broth (TB) medium with the required antibiotic.
- Wells get inoculated with a loop of the cryoculture or an single colony from an agar plate and grown over night at 37°C, 200 rpm sealed with a breathable membrane.
- After the overnight, 3 μL of each preculture are transferred to fresh 300 μL TB + antibiotic and grown for 3 h, 37°C, 200 rpm. Then cultures get induced with 1 mM IPTG (or other appropriated inducer) and grown over night at 30°C, 200 rpm.
- Then, the cultures are spin down in the plate, the media is removed and plates are covered with a fresh breathable membrane and frozen for at least 6 h at -30°C.
- Finally the plates get lyophilized overnight.
- For assays, the liophilized cells are re-suspended in 0.6 ml of a buffer with low ionic strenght and close to neutral pH; 2-10 μL are used for activity tests.

## 5. Protocols relevant for TEXTILE APPLICATIONS

As detailed in Deliverable D2.1, priority enzymes are those for textile fibres, that include cotton (CO), polyester fibres (PES), polyamide fibres (PA), and elastane (polyether-polyurea copolymer): Lipases and

polymer-degrading enzymes (herein referred as polyesterases), cutinases, amidases, proteases (subtilisin, papain, alcalase and bromelain-type), polyurethanases, cellulases, amylases and bleaching and dye decolorizing enzymes (oxidoreductases). Among them, priority targets are lipases for removing residual spinning oils/sizing products that, if not eliminated, will otherwise generate emissions during the drying and fixation steps; priority textiles are those made of polyester (PES). Priority targets are also oxidoreductases (laccase-like) for supporting in the decolorization of dyes. Currently, these additives/preparation materials and residual dyes are removed by a water/surfactant process and reducing the rinsing steps/duration is the expected goal when adding enzymes into the cleaning processes, and enzymes are requested to support the elimination of those additives/preparation materials and residual dyes in water. Final priority target are polyesterases that can be applied in the biodegradation of the current textile materials in such a way that they can even be reused to produce new recycled textiles.

The following decision-taken strategy, once activity towards lipase and dye-related substrates is confirmed, has been undertaken.

#### Decision taken screen 1:

Enzymes are capable of eliminating spinning oils/sizing products from raw fabrics provided by
partner Schoeller, in buffer or water.
Note: As a control the pretreated fabrics from which the spinning oils / sizing products have been
chemically and thermally removed, are used. No control enzyme can be used, as no lipases are
used commercially for this purpose.

#### Decision taken screen 2:

• Enzymes are capable of eliminating dye products provided by partner Schoeller, in buffer or water. Note: No control enzyme can be used, as no lipases are used commercially for this purpose.

#### Decision taken screen 3:

 Enzymes are capable of degrading polyester (PES) fabrics provided by partner Schoeller, in buffer or water.
 Note: As a control, best performing PETases reported in the literature are used.

Note: As a control, best performing PElases reported in the literature are used.

Below, the fabrics and dyes provided by partner Schoeller, and that were used as real substrates for activity tests when appropriated, are detailed:

#### Raw and pre-treated fabrics (to remove the spinning oils, sizing products, etc.)

Sent from Schoeller to CSIC, and CSIC distributed them to Bangor, IST-ID, UDUS, UHAM, and BioC\_Chem (September 2021).

Variant	Туре	Article No.	Based material	Available Status in stock	Comp. <sup>1</sup> / Weight
1	Woven	61488	61488Z	Raw	92% PA, 8% EL 180 g/m <sup>2</sup>
			61488Z	Pre-treated	
2	Woven	61988	61988F1	Raw	92% PA, 8% EL 280 g/m <sup>2</sup>
			61988F1	Pre-treated	
3	Woven	67007	67007	Raw	88% PA,12% EL 135 g/m <sup>2</sup>
			67007	Pre-treated	

5 Woven 66299 5237/00 Raw 92% CO, 8% EL	g/m²
	. 240 g/m <sup>2</sup>
6 Warp-knitted E03130 E03130 Raw 80% PA, 20% EI	L 160 g/m <sup>2</sup>

<sup>1</sup>PA: Polyamide; EL: Elastane; PES: Polyester.

Sample¤	Article Nr.¤	Condition¤	A.→Quantitative· [%]*¤	B. Qualitativ [IR-Spectrum]∞
1¤	3X58¤	raw¤	1.6¤	Fatty-acid-ester-or-emulsified¶ Mineral-oil-/-paraffin¤
		Pretreated¤	0.0¤	Quantity-too-small-for-analysis <sup>q</sup>
2¤ E03130¤		raw¤	1.9¤	Fatty-acid-ethoxylate,-amide,-siliconeg
29	E03130¤	Pretreated¤	0.1¤	Quantity-too-small-for-analysis <sup>q</sup>
3¤	61988F1¤	raw¤	0.7¤	Fatty-acid-ester-(-EO),amide, -little-siliconeg
20	01900F1¤	Pretreated¤	0.1¤	Quantity-too-small-for-analysis <sup>q</sup>
4g 5237-00¤		raw¤	0.7¤	Fatty-acid-esters,-silicone¤
49	5257-00 <sup>10</sup>	Pretreated¤	0.5¤	Fatty-acid-esters,-silicone¤
5ø	61488F1¤	raw¤	2.6¤	Fatty-acid-ethoxylate, amide-(little-amount)¤
50 01400F10		Pretreated¤	0.2¤	Fatty-acid-ester, polyamide, siliconeg
6¤	67007¤	raw¤	2.5¤	Mineral-oil-/-paraffin,-fatty-acid-ethoxylate,- amide,-silicone¤
6 <u>2</u> 6/00/¤		Pretreated¤	1.2¤	Fatty-acid-esters,-silicone,-possibly-mineral-oil- /-paraffin¤





#### Dyeing baths and chemicals inside (to remove the dyes from it - decolorization)

Sent from Schoeller to CSIC, Bangor, IST-ID, Inofea (received on 15.11.2022)

Sent from CSIC to UDUS and (received on 24.11.2022)

Sent from Inofea to FHNW (received on 24.11.2022)

Schoeller Receipt Nr. 31964900

Proceedings (Verfahren): V-AN4SDNFPH4RFDI

Dyeing of materials on Noseda machinery with pH 4.5

Chemicals in the dyeing bath:

Chemicals	Amount (g/L)	Provider
Periwet WDP NEW	0.4	Petry Chemie
BIAVIN BPA	2	СНТ
SARABID C14	0.65	СНТ
Ammoniumsulfate-LSG. 33%	4	Bilgram Chemie GmbH
Ameisesäure 85%	0.6	Bilgram Chemie GmbH
Dyeing material		
BEMAPLEX SCHWARZ D.HF	3.57	СНТ



## 5.1. Agar plate polyesterase screening assay with PET-nanoparticles

## 5.1.1. Protocol for the preparation of PET-nanoparticle solution

- Dissolve completely 0.1 g polyethylene terephthalate (PET)<sup>1</sup> in 10 mL 1,1,1,3,3,3-hexafluor-2-propanole (Sigma-Aldrich/Merck) and pour it in a burette.
- Slowly transfer this solution (drop-by-drop) into a 100 mL ice-cold water containing beaker (in an ice bath), which is strongly agitated, to produce precipitated PET polymer.
- Remove the solvent<sup>1</sup> and part of the water by distillation, so that the 110 mL total volume is reduced to approx. 2-fold (approx. 50 mL final volume).
- The concentrated water solution containing precipitated PET-nanoparticles can be stored at 4°C for about 2 months. Usually we get a solution of about 0.8-1.6 mg/ml nanoparticles in water.

<sup>1</sup>Any kind of plastic bottle can be used as source of PET. We used PET from fruit juice bottle of *Granini*.

## 5.1.2. Preparation of PET-nanoparticle screening plates

- Mix 50 mL medium (M9 minimal medium plus 1.8% (w/v) agar agar)<sup>2</sup> with the concentrated nanoparticle solution (~12 mL).
- Those plates can be stored for a maximum of 2 weeks at 4°C.
- Place bacterial<sup>3</sup> or protein<sup>4</sup> solution onto the plate.
- Usually we observe a lysis zone<sup>5</sup> around the colonies after 2-3 days of incubation at 37°C, depending on the activity of the enzymes.

<sup>1</sup>It is absolutely mandatory to completely remove the solvent prior to the incorporation of the nanoparticles into the plates (see below), otherwise the bacteria and enzymes are strongly growth-inhibited.

<sup>2</sup>This can be adapted to any other buffer and medium with appropriate antibiotic.

<sup>3</sup>Grow *Escherichia coli* expressing the enzyme of interest at 37°C on Luria Bertani (LB) agar medium, supplemented with an appropriate antibiotic. Pick a single colony to inoculate 1 mL of LB-antibiotic medium in a 2-mL Eppendorf and cultivate it at 37°C and 150 rpm during 6 h ( $OD_{600nm}$  ranging from 0.7 to 0.9). Afterwards, use 5  $\mu$ L of this culture to inoculate the PET-nanoparticle plate, and incubate it at 37°C.

<sup>4</sup>If purified or cell extract of protein is available, use 5 μL of a concentrated protein solution to inoculate the PETnanoparticle plate, and incubate it at the desired temperature.

<sup>5</sup>The diameter of the lysis zone is directly correlated to the hydrolytic enzyme activity; we observed diameters ranging from mm to 1 cm scale (low activity:  $\emptyset$  < 1 mm; medium activity:  $\emptyset$  1-3 mm; high activity:  $\emptyset$  >3 mm).

## 5.2. Agar plate polyesterase screening assay with polylactic acid (PLA) emulsion

- To prepare the PLA (poly(DL)-lactide, average Mw 2000 or higher, purchased from PolySciTech) emulsion (Teeraphatpornchai T. *et al.*, 2003, Biotechnol. Lett., 25: 23-28), dissolve 2 g of analytical grade PLA in 40 mL of dichloromethane, mixed with 1 L of 50 mM Tris-HCl pH 8.0 containing 0.1 g of Triton X-100 as surfactant and subject it to sonication for 10 min.
- Heat the stable emulsion at 80°C for 2 h to remove the solvent; then mix it with 15 g of agarose or agar (1.5% final agarose, 2% final PLA) and pour it into Petri dishes.
- Cool to room temperature.
- Punch manually the well in the solidified agarose and load it with purified proteins (50 µL, 10-50 µg/well).
- Seal the plates and incubate them at 30°C for 1-3 days. The presence of PLA degrading activity is indicated by the formation of a clear halo around the wells. To compare polyesterase depolymerization activity of different enzymes, image analysis is conducted using the histogram tool on Adobe Photoshop software, and pixel counts are plotted on the graph with error bars representing measurements with varying pixel tolerance values.

## 5.3. Agar plate polyesterase screening assay with PET or PLA polymers

The protocol is based on Hajighasemi et al. (2016, 2018) and can be used with PET and PLA polymers as substrates.

- Dissolve 0.2% (or enough polymer to get a good contrast to visualise the halos) of polymer in 1-2 mL of solvent (acetone, chloroform or dichloromethane) (*3PET dichloromethane*, *PLA acetone/chroloform*)<sup>1</sup>.
- Dissolve 1-1.2% agarose in 50 mM Tris-HCl pH 8.0. Microwave to dissolve agarose/agar and then keep solution at 50-60°C stirring. Add 0.1-0.01% Plysurf A210G (a surfactant; Daiichi Kogyo Seiyaku, Tokyo, Japan).
- Add solvent solution drop by drop in the aqueous solution while vigorously stirring.
- Sonicate at maximum power.
- Heat (stirring, under fume hood) at 80°C to remove the solvent.
- Pour in Petri dishes.
- Hydrolysis is observed by the appearance of a lysis zone.

<sup>1</sup>Alternatively, prepare 2x solution of agarose/agar and 2x solution of aqueous phase + solvent and mix after evaporation of solvent.

### 5.4. Agar plate polyesterase screening assay with Impranil DLN

- Add 0.5% Impranil DLN-SD emulsion (COVESTRO, Leverkusen, Germany) per 1 L of sterile molten LB agar and mix thoroughly, e.g. using an Ultra Turrax.
- Pour 25 mL medium into appropriate Petri dishes and allow the agar to solidify for at least 15 min.
- Plate bacterial clones or specific microorganism at optimal growth temperature for at least 16 h.
- Hydrolysis is observed by the appearance of a lysis zone.

## 5.5. Agar plate polyesterase screening assay with polycaprolactone diol (PCDMn530)

- Prepare a 50% (v/v) PCD (average Mn 530 Da) emulsion: mix the PCDMn530 (Sigma-Aldrich/Merck) and 50 g/L arabic gum with sterile distilled water. Homogenize the mixture for at least 1 min to yield a stable emulsion, e.g. using an Ultra Turrax<sup>1</sup>.
- Add 30 mL of PCDMn530 emulsion per 1 L of LB agar and mix thoroughly, e.g. using an Ultra Turrax.
- Pour 25 mL medium portions into appropriate Petri dishes and allow the agar to solidify for at least 15 min.
- Plate bacterial clones or specific microorganism and incubate at optimal growth for at least 16 h.
- Hydrolysis is observed by the appearance of a lysis zone.

<sup>1</sup>Add the arabic gum powder to the respective volume of water. Filling water into a tube or a bottle with a layer of the powder at the bottom should be avoided because it will result in a hard-to-dissolve clot of gum.

#### 5.6. Agar plate polyesterase screening assay with polycaprolactone (PCL) nanoparticles

The protocol derived from Jarrett et al. (1984).

- Prepare a 5 g/L PCL solution by completely solving PCL (average Mn ~10 000 by GPC, density 1.146 g/mL, Sigma-Aldrich/Merck) in pre-heated acetone at 50°C under continuous stirring. Pre-heat an appropriate volume of sterile water likewise to 50°C for the next step<sup>1</sup>.
- Prepare a PCL particle suspension by slowly pouring the PCL solution drop by drop under continuous stirring into the water until a final acetone percentage of aprox. 10-15% is reached.

- Add 100 mL of the warm PCL suspension per 1 L of LB agar and mix thoroughly, e.g. using an Ultra Turrax.
- Pour 25 mL medium into appropriate Petri dishes and allow the agar to solidify for at least 15 min.
- Plate bacterial clones or specific microorganism and incubate at optimal growth temperature for at least 16 h.
- Hydrolysis is observed by the appearance of a lysis zone.

<sup>1</sup>A turbid dispersion should be formed. Pour carefully, because too fast supplementation of PCL solution easily leads to the formation of tiny globular plastic particles instead of a homogenous suspension.

## 5.7. Agar plate protease skim milk screening assay

- Autoclave 3% skim milk in water and LB-Medium containing 3% Agar-Agar separately.
- Mix the above two solutions in a ratio of 1:1.
- Plate bacterial clone or specific microorganism<sup>1</sup> and incubate at optimal growth temperature for at least 16 h.
- Hydrolysis is observed by the appearance of a lysis zone.

 $^1$  It is recommended to place into the plates a volume of 10  $\mu L$  of a pre-grown culture.

### 5.8. Agar plate protease casein screening assay

- Prepare a calcium caseinate agar plate following the protocol by Sigma-Aldrich/Merck (ref. 21065-500G). Autoclave.
- Plate bacterial clone or specific microorganism<sup>1</sup> and incubate at optimal growth temperature for at least 16 h.
- Hydrolysis is observed by the appearance of a lysis zone.

<sup>1</sup>It is recommended to place into the plates a volume of 10  $\mu$ L of a pre-grown culture.

#### 5.9. Agar plate cellulase screening assay

This protocol is based on the use of the  $\beta$ -glucose polymer Avicel, and on the ability of the exo  $\beta$ -glucosidases to hydrolyze the polymer. That hydrolysis results in a clear halo formation surrounding colonies.

- Carry out the screening is carried out on solid medium containing 1.0% w/v Avicel (Sigma-Aldrich/Merck) partially hydrolysed<sup>1</sup>. Then, maintain the plates up to 16-48 h at 37°C. Positive activity is detected by the presence of a halo.
- Slowly add 10 mL of ice-cold 86.2% H<sub>3</sub>PO<sub>4</sub> to the slurry with vigorous stirring so that the final phosphoric acid concentration is approx. 83.2%. Before adding the last 2 mL of phosphoric acid, mix evenly the cellulose suspension solution. The cellulose mixture turns transparent within several minutes, and stands for approx. an hour on ice with occasional stirring.
- Add approx. 40 mL of ice-cold water at a rate of around 10 mL per addition with vigorous stirring between additions, resulting in a white cloudy precipitate.
- Centrifuge the precipitated cellulose at ~5000 g and 4°C for 20 min. Resuspend the pellet with ice-cold water, followed by centrifugation to remove the supernatant containing phosphoric acid four times. Add approx. 0.5 mL of 2 M Na<sub>2</sub>CO<sub>3</sub> to neutralize the residual phosphoric acid, and then, use 45 mL of ice-cold distilled water to suspend the cellulose pellet.
- After centrifugation, resuspend the pellet in distilled water and centrifuge twice or until pH 5.0-7.0.

<sup>1</sup>Hydrolyzed Avicel is prepared as follows (adapted from Biomacromolecule 2006 Feb;7(2):644-8): Approximately 0.2 g of microcrystalline cellulose is added to a 50 mL centrifuge tube, and 0.6 mL distilled water is added to wet the cellulose powder to form a cellulose-suspended slurry. The regenerated amorphous (homogeneous) cellulose slurry can be kept for a long time at ~4°C, and used for activity screening at 1% w/v (see above).

## 5.10. Agar plate oxidoreductase (laccase/oxidase) screening assay

In order to screen purified proteins for oxidase activity against polymeric lignin, we designed a lignin-agarose plate screen, which is based on a polyester-agarose screen developed for screening purified proteins for polyesterase activity (Hajighasemi M. et al., 2016, Biomacromolecules, 17: 2027-2039):

- Load purified proteins (up to 50 μg) into small wells in 1.5% agarose containing low sulfonate kraft lignin or Indulin AT kraft lignin (0.025-0.25%) in 100 mM Tris-HCl buffer pH 7.0.
- Incubate overnight at 37°C.
- Stain the lignin-agarose plates with a mix (50/50) of 1% (w/v) potassium ferricyanide (K<sub>3</sub>[Fe(CN)<sub>6</sub>]) and 1% ferric chloride (FeCl<sub>3</sub>) (a staining protocol developed for fungi (Mswaka A.Y., Magan N., 1998, Mycol. Res., 102: 1399-1404; Peterson R.A. et al., 2009, Lett. Appl. Microbiol., 48: 218-225), resulting in the formation of a green background. The development of green background involves two reactions: (1), generation of Fe<sup>2+</sup> (from the oxidation of hydroxyl groups of lignin phenolic units by FeCl<sub>3</sub>), which (2) reacts with K<sub>3</sub>[Fe(CN)<sub>6</sub>] generating green colored Prussian blue (KFe<sub>3</sub><sup>+</sup>[Fe<sup>2+</sup>(CN)<sub>6</sub>]). The presence of lignin oxidase activity is detected as the formation of yellow halos around positive wells (due to the decrease in the content of phenolic-hydroxyl groups).
- Yellow halo areas are quantified in pixels obtained via image scanning as described previously (Hajighasemi M. et al., 2018, Environ. Sci. Technol., 52: 12388-12401).

# 5.11. Liquid protocol for quantifying esterase-lipase activity with chromogenic commercial kits

## 5.11.1. NEFA-HR(2) kit

In the first steps of a textile manufacture, the spinning oils used to produce the continuous thread or the yarn need to be eliminated. Traditionally, this step uses huge amounts of water. The oils can be removed by the action of enzymes to replace or highly reduce the water usage. To test this, the NEFA-HR(2) kit (Waco, 517982 - Reagent 1 Set kit NEFA-HR(2) Assay, 517984 - Reagent 2 Set, kit NEFA-HR(2) Assay, 517954 - Standard, kit NEFA-HR(2) Assay), also referred to as ACS-ACOD method, can be is used (quantitative determination of Non-Esterified Fatty Acids - NEFA). NEFA in sample is converted to Acyl-CoA, adenosine monophosphate (AMP) and pyrophosphoric acid (PPi) by the action of Acyl-CoA synthetase (ACS), under coexistence with coenzyme A (CoA) and adenosine 5-triphosphate disodium salt (ATP). Obtained Acyl-CoA is oxidized and yields 2,3-trans-Enoyl-CoA and hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) by the action of Acyl-CoA oxidase (ACOD). In the presence of a peroxidase, the H<sub>2</sub>O<sub>2</sub> formed yields a blue-purple pigment by quantitative oxidation condensation with 3-Methyl-N-Ethyl-N-( $\beta$ -Hydroxyethyl)-Aniline (MEHA) and 4-aminoantipyrine (4-AA). Non-esterified fatty acids (NEFA) concentration is obtained by measuring absorbance of the blue-purple color.

Reagent preparation:

- R1: Prepare R1 by mixing one bottle of Color A and Solvent A. After preparing the R1, store at 2 10°C and use within 1 month.
- R2: Prepare R2 by mixing one bottle of Color B and Solvent B. After preparing the R2, store at 2 10°C and use within 1 month.




\_

Protocol for spinning oils removal from yarn (adapted from the kit specifications):

- Incubate the enzyme (at different concentrations) with a small piece (4 mg) of the yarn material (e.g. provided by Schoeller Textile), previoulsy cutted using a circle hole puncher (5 mm diameter), in 100 μL of 40 mM of the appropriated buffer (e.g. pH 7.0 HEPES buffer) for 24h, at 30°C and 950 rpm.
  - The reaction outcome is tested using the NEFA-Kit, following specifications:
    - Transfer 10  $\mu L$  of the reaction solution + 100  $\mu L$  of NEFA solution 1 to a 96 microplate well. Incubate 6 min at 37°C.
    - Add 50  $\mu$ L of NEFA solution 2. Incubate 6 min at 30°C.
    - Measure sample's absorbance at 550 nm using a microplate spectrometer (e.g. Synergy HT Multi Mode Microplate Reader, Agilent, Madrid, Spain).

A summary of the protocol is detailed, in a schematic manner, below:



## 5.12. Liquid polyesterase quantification assay with polylactic acid (PLA) powder

- To determine PLA depolymerase activity against solid PLA, incubate small amounts of PLA powder (10-15 mg) in a reaction mixture (1 mL) containing 400 mM Tris-HCl buffer pH 8.0.
- Then, add 50 μg of protein with shaking at 35°C for 36 h.
- At different time intervals, take small samples and clarify them using centrifugal filters (Mw cut-off 10 kDa); measure free lactate using L- and/or D-lactate dehydrogenase (LDH) added in excess (0.5 mg/mL) which enables the detection of both D- and L-enantiomers of lactic acid (Babson A.L., Phillips G.A., 1965, Clin. Chim. Acta, 12: 210-215). The oligomeric PLA products in flow-through aliquots (90 μL) were converted to monomeric lactic acid after 5 min incubation at 95°C with 10 μL of 10 mM NaOH followed by LDH assay.

### 5.13. Liquid polyesterase quantification assay using Impranil DLN-SD

- Prepare a substrate solution by mixing 0.5 % (vol/vol) Impranil DLN-SD dispersion in 100 mM KPi pH 7.2.
- Pipette 10 μL of enzyme solution into each well of a 96-well plate (10 μL enzyme buffer as a control).
- Add 190 µL of substrate solution.
- Follow the clearing of the reaction (turbidity clearing) by measuring the absorbance at 580 nm.

## 5.14. Liquid protocol for the quantification of BHET or PET foil hydrolysis using UHPLC

- Prepare the enzyme reaction in 1.5 mL Eppendorf tubes. The reaction mixture is composed of 500 nM purified enzyme in 20 mM potassium phosphate buffer pH 7.4 with 20 % (v/v) DMSO in a total volume of 300  $\mu$ L. As a substrate, add either 0.75  $\mu$ L of 400 mM bis(2-hydroxyethyl) terephthalate (BHET; 95% purity, Sigma Aldrich) dissolved in DMSO or a circular piece of PET film (6 mm diameter) to each tube. A reaction without the addition of an enzyme is used as a control<sup>1</sup>.
- Incubate the reaction for 48 h at 30°C, 800 rpm on a shaker.
- Stop the reaction by heat inactivation of the enzymes (e.g. 10 min at 80°C) or separate reaction products and enzymes by ultrafiltration (e.g. centrifugal filters with 10 kDa molecular weight cut-off (MWCO)).
- Quantify reaction products terephthalic acid (TPA), mono-(2-hydroxyethyl) terephthalate (MHET), and bis-(2-hydroxyethyl) terephthalate (BHET) via HPLC using pure standards.
- HPLC setup:

Acquity UPLC (Waters GmbH) equipped with an Acquity UPLC BEH C18 column (1.7 mm particle size). HPLC program:

Mobile phase (A) 20 mM Na<sub>2</sub>HPO<sub>4</sub> pH 2.5 (pH adjusted with H<sub>2</sub>SO<sub>4</sub>) and (B) methanol, the effluent was monitored at 240 nm. Column temperature set to 35°C. Flow rate 0.208 mL/min. Injection volume 4  $\mu$ L. 75% (A) and 25% (B) for 1.28 min, followed by a linear gradient to 100% (B) in 2 min, hold 100% (B) for 3 min, linear gradient from 100 to 25% (B) in 1 min and hold 25% (B) until minute 8.28 is reached<sup>2</sup>.

<sup>1</sup>For standardized PET film substrate we recommend: (A) Polyethylene terephthalate Film ES301445, Thickness: 0.25 mm, Transparent, Amorphous, Goodfellow GmbH, order no. 518-097-996. (B) Polyethylene terephthalate Film ES301450, Thickness: 0.25 mm, Biaxially Oriented, Clear, Goodfellow GmbH, order no. 543-716-953.

<sup>2</sup>For MHET no commercial standard is available. A standard curve can be calculated from a series of hydrolysis reactions with BHET as a substrate and the quantified amounts of BHET and TPA.

## 5.15. Liquid oxidoreductase (laccase/oxidase) quantification assay

- Prepare reaction mixtures (200 μL) containing 100 mM Britton-Robinson buffer, optimal pH for each substrate (pH 2.0 9.0, see below), substrates (0.04 20 mM)
- Add 10 µg of enzyme (for preliminary screening, then adjust to 0.1-0.5 µg of protein).

Incubate at 30°C in 96-well plates and measure the conversion at the appropriated wavelenght to each of the selected substrates.
2,2'-azinobis(3-ethylbenzthiazoline-6-sulfonate) (ABTS), pH 5.0 (ε = 36 mM<sup>-1</sup>cm<sup>-1</sup>, 420 nm) Syringol, pH 7.0 (ε = 14.8 mM<sup>-1</sup>cm<sup>-1</sup>, 468 nm) Guaiacol, pH 7.0 (ε = 6.4 mM<sup>-1</sup>cm<sup>-1</sup>, 436 nm) *p*-phenylenediamine, pH 6.0 (ε = 14.7 nM<sup>-1</sup>cm<sup>-1</sup>, 487 nm) 3-hydroxyanthranilic acid (3-HAA), pH 6.0 (480 nm) Gallic acid, pH 6.0 (480 nm) Sinapic acid, pH 8.0 (480 nm)

## 5.16. Liquid amylase quantification assay with chromogenic substrates

- Prepare reaction mixture (200 μL) containing 50 mM buffer (HEPES-K pH 7.0; MOPS-K pH 6.5; MES-K pH 6.0), 1 mM substrate (*p*-nitrophenyl α-D-maltoside or β-D-maltoside, available from Sigma/Merck), 20 mM NaCl, 1 mM DTT, and 1-5 µg of purified protein.
- Measure substrate hydrolysis spectrophotometrically after 10 min incubation at 30°C (or room temperature) at 420 nm in a microplate reader.

## 5.17. Liquid amylase quantification assay with polymeric substrates

- Add 40  $\mu L$  of starch (2 mg/mL) to 40  $\mu L$  of 0.1 M HEPES-K buffer pH 7.0 in a microplate covered with a plastic film.
- Incubate the microplate for 30 min at 30-50°C.
- Stop the reaction by adding 20  $\mu L$  of 1 M HCl followed by the addition of 100  $\mu L$  of iodine reagent (5 mM  $I_2$  and 5 mM Kl).
- Measure activity by following color development at 580 nm using a plate reader (Xiao Z. et al., 2006, Anal. Biochem., 351: 146-148).

An alternative method consists in the detection of reducing sugars with a modified 3,5-dinitrosalicylic (DNS) method adapted to a 96-well microplate scale.

- Prepare the DNS solution: 10 g 3,5-dinitrosalicylic acid (DNS), 300 g sodium and potassium tartrate (Rochelle salt), 16 g NaOH and distilled water up to 1 L (*Note: it takes some time until completely dissolved*).
- In a microtiter plate or 1.5 mL Eppendorf, add 45 μL buffer, 5 μL reaction solution (see above) and 50 μL DNS solution.
- Incubate at 85°C for 25 min (in these conditions, the reducing sugars allow the transformation of DNS into DAS (3-amino-5-nitrosalicylic acid) with the color change from yellow to dark brown).
- Add 100 μL water and read the absorbance at 540 nm.

## 5.18. Liquid proteolytic activity assays using FITC-labelled protein

- Prepare labelled protein stock in water (5 mg/mL), using Thermo Scientific Kit or self-made Fluorescein isothiocyanate (FITC) labelled protein.
- Prepare working reagent (WR) of 100 μg/mL labelled protein in Tris Buffered Saline (TBS) Buffer.
- Add 100 μL sample (or standard) to each well of a 96-well plate (use 37.5 μL/well for 384-well plate). In addition to the zero-standard, prepare a blank using a buffer similar to that used for the protease sample.
- Add 100 μL of WR (1:500) to all wells of a 96-well plate (use 37.5 μL/well for 384-well plate).
- Incubate for 5-60 min at room temperature.
- Measure fluorescence in a plate reader using a fluorescein excitation/emission filter set.

- Subtract the blank from each sample and standard measurement and then prepare a standard curve: for fluorescence resonance energy transfer (FRET), plot the change in relative fluorescence units (RFU) of the standards vs. protease concentration.

## 5.19. Liquid proteolytic (alcalase) activity assays using *p*-nitroanalide chromophores

Alcalase-like activity is measured in batch through a fluorometric assay, using a synthetic specific substrate (N-succinyl-l-alanyl-l-alanyl-l-ananyl-*p*-nitroanilide; available from Merck/Sigma). Incubate protein samples (purified protein, cell extract or cell suspensions) with a buffer solution (150 nM Tris-HCl pH 7.5, 200 nM NaCl, 10 uM ZnCl<sub>2</sub>) and the substrate (1 mg/mg stock solution in DMSO) (portions 10:99:1, respectively). After incubation at desired temperature for up to 16 h, quantify the absorbance at 405 nm with a microtiter plate reader.

## 5.20. Liquid proteolytic (subtilisin) activity assays using *p*-nitroanalide chromophores

Alcalase-like activity is measured in batch through a fluorometric assay, using a synthetic specific substrate (N-succinyl-Ala-Ala-Pro-Phe-*p*-nitroanilide; available from Merck/Sigma). Protein samples (purified protein, cell extract or cell suspensions) are incubated with a buffer solution (150 nM Tris-HCl pH 7.5, 200 nM NaCl, 10  $\mu$ M ZnCl<sub>2</sub>) and the substrate (1 mg/mg stock solution in DMSO) (portions 10:99:1, respectively). After incubation at desired temperature for up to 16 h, the absorbance at 405 nm is quantified with a microtiter plate reader.

## 5.21. Liquid proteolytic activity assays using azocasein

- Mix 1.1 mL of 50 mM sodium phosphate buffer pH 7.6 (standard buffer) containing 10 μg of protein (or appropriated amount of cell extract or cell suspension) with 0.65% (w/v) of the substrate azocasein.
- After incubating the reaction mixture at 30°C for 30 min, add 900  $\mu$ L of 110 mM trichloroacetic acid and incubate the mixture at 37°C for 30 min.
- To remove precipitant, centrifuge the mixture (12,000 g for 3 min at 4°C) and filter through a 0.45-µm filter.
- Mix the filtrate (500  $\mu$ L) with 250  $\mu$ L Folin & Ciocalteu's phenol reagent and 1.25 mL of 500 mM sodium carbonate solution.
- After 30 min at 37°C, determine the  $OD_{660}$  and calculate the enzyme activity based on an L-tyrosine standard curve.
- One unit of protease activity is defined as the activity required to produce 1 nmol of tyrosine equivalents released amino acids per min, per mg of protein at 30°C.

## 5.22. Liquid hydrophilic and dyeability modification of fabrics

Protocol adapted from Gao et al 2017 (Aiqin Gao, Hongwei Shen, Hongjuan Zhang, Guanchen Feng, Kongliang Xie, Journal of Cleaner Production 2017, 164, 277-287). This protocol can be used for proteases, lipases, cutinases and polyesterases.

- Prepare a buffer aqueous solution containing certain concentrations of lipase.
- Immerse the polyester, PLA, PET or polyamide fabric (5.0 g) in the solution (100 mL).
- Perform the treatment at the desired temperature and pH 7.5 for a certain time.
- After the enzymolysis reaction, increase the temperature to 100°C, hold for 15 min, and then cool, to inactivate the lipase.
- Determine the polyester hydrolysis degree from the quantities of terephthalic acid (TPA) derivatives (TPA, ethylene terephthalate, and mono(2-hydroxyethyl) terephthalate) released into the samples supernatant by measuring the absorbance at 241 nm. The possible hydrolysis products were TPA, ethylene

terephthalate, and mono(2-hydroxyethyl) terephthalate. Use a TPA calibration curve to calculate the total amount of products released (all three products have the same molar absorption, arising from their carbonyl groups).

Calculate the activity by applyign the following equation:

$$A = -6.4690 \times 10^{-4} + 75.1195 \times C$$

where A is the absorbance at 241 nm and C is concentration of TPA (mg/mL).

Hydrolytic activity on polyester, PLA, PET or polyamide fabrics can be also measured by a titration method (protocol adapted from H.R. Kim and W.S Song (Fibers and Polymers 2010, 11, 67-71). This protocol can be used for proteases, lipase, cutinase, polyesterases.

- Cut each fabric sample into 4×4 cm pieces that weight approx. 0.1 g.
- Place each sample into a 20 mL vial bottle containing 8 mL of an appropriate buffer. Then, treat the polyester fabrics with different temperature, pH, treatment time, and concentrations of enzymes. All treatments are performed at 150 rpm using a shaking waterbath.
- After the treatments, transfer the test solutions to a 50 L Erlenmeyer flask, and add 20 mL ethanol to the test solution to inactivate its enzymes. Add four drops of 0.9% TPH indicator to the test solution. The test solution is titrated with 0.1 M NaOH into a light blue color. Then, record the volume of the 0.1 M NaOH used in the sample test.

Enzymatic hydrolytic activity can be also measured according to the method of SH Lee and SY Yeo (Fibers and Polymers 2016, 17, 1154-1161). This protocol can be used for proteases, lipase, cutinase, polyesterases.

- Mix a PLA fabric or other polymer-based fabric (0.5 g) with 50 mL of 0.1 M sodium phosphate buffer pH
   7.0. Then, add 2 mM *p*-NP<sup>1</sup> as a yellow chromogen pH indicator.
- Place each sample into a 20 mL vial bottle containing 8 mL of an the above prepared solution.
- Then, treat the fabrics with different temperature, pH, treatment time, and concentrations of enzymes. All treatments are performed at 150 rpm using a shaking waterbath.
- Measure the hydrolytic activity by taken aliquots and following the absorbance at 405 nm. Calculate the hydrolytic activity of the enzyme on the basis of absorbance at 405 nm using the following equation: Hydrolytic activity = A405 after hydrolysis – A405 before hydrolysis.

 $^{1}p$ -nitrophenol (*p*-NP) could be substituted by 0.45 Phenol Red, and in this case, the buffer 100 mM sodium phosphate buffer(pH 7.0, need to be substituted by 5 mM EPPS buffer.

## 5.23. Liquid amino modification of polyamide fabrics

Protocol adapted from Kanelli et al (Maria Kanelli, Sozon Vasilakos, Spyridon Ladas, Emmanouil Symianakis, Paul Christakopoulos, Evangelos Topakas, Process Biochemistry 2017, 59, 97-103). This protocol can be used for proteases, amidases and lipases. The hydrolysis of polyamide model occurred as followed:

- Treat 50 mg of the model fabric with an appropriated amount of enzyme in 1.5 mL potassium phosphate buffer pH 8.0 and at appropriated temperature, overnight.
- Subsequently, filter the samples to obtain the soluble fraction of the hydrolysis and analyze them with the Trinitrobenzenesulfonic acid (TNBS) method. This method is based on the reaction of the primary amino groups with the sodium salt of TNBS leading to the formation of a complex that can be measured spectrophotometrically at 420 nm. Briefly, incubate 1 mL of the reaction's supernatant with 25 μL of 30 mM aqueous TNBS for 30 min at 30°C. For the quantification of amino-groups that is indicative of the enzymatic hydrolysis of the PA model, the absorbance is read against the controls, one with model fabrics

and buffer and one with enzyme and buffer at 420 nm in a spectrophotometer. Prepare a calibration curve using hexamethylenediamine as a standard solution in different concentrations.


## 5.24. Dye liquid discoloration test

- Prepare the reaction mixture in 96-well microplates by adding 20  $\mu$ L of concentrated dye (provided by partner Schoeller), 10  $\mu$ L of 2 mM mediator (ABTS) (for peroxidase screen prepare fresh 0.06% H<sub>2</sub>O<sub>2</sub> add 100 uL to the reaction mixture), and water to 200  $\mu$ L taking into account the enzyme volume (5-10  $\mu$ g).
- (Control wells should a) not contain enzyme, but the elution buffer, b) contain enzyme and mediator, but no dye).
- Mix by pipetting (plate shaker alternatively for 10 min on 500 rpm).
- Add 5-10 μg of purified protein to fit in the final volume 200 μL.
- Mix by pipetting and read the 0 time point at 588 nm , save the data.
- Seal plate with adhesive seal and keep shaking for overnight at 30°C, 500 rpm.
- Next day shortly spin down the plate and read at 588 nm.
- Subtract the ON data from the 0 time point read, taking into account the controls.

Notes:

For futher protocol optimization reaction volume may be considered to be decreased to 150  $\mu$ L for better aeration and the concentration of mediatior may need optimization.

As shown in the Figure below, 588 nm was identified as maximal absorbance for the dying liquid cocktail from Schoeller, dye absorbance is distant from the ABTS absorbance (420 nm), if using different mediators, their extinction should be taken into account.




# 6. Protocols relevant for COSMETIC APPLICATIONS

As detailed in Deliverable D2.1, priority enzymes are those for processing hyaluronic acid:

- Heparanase (EC 3.2.1.166)
- Hyaluronate lyase (cd01083 EC 4.2.2.1)
- Hyaluronidase (EC 3.2.1.35, EC3.2.1.36, pfam03662, pfam01630)

The idea is to identify hyaluronidase-like enzymes capable of degrading hyaluronic acid to small hyaluronic acid products with 1-2 k Dalton molecular weight (see figure below), at <37°C, no solvents, and high viscosity solutions.



The following two real hyaluronic acid products were provided by partner Evonik for activity tests:

- Low molecular weight (50 kDa) hyaluronic acid HyaCare<sup>®</sup> 50
- Low molecular weight hyaluronic acid (<10 kDa), Hyalo-Oligo

This material was provided to CSIC directly by Evonik, and then distributed to Bangor, UDUS, UHAM, IST-ID, and CNR (September 2021).



#### 6.1. Liquid colorimetric assay for hyaluronidase activity

The hyaluronidase activity is determined by detection of reducing sugars with three different colorimetric methods. We may use hyaluronic acid, sourced from Sigma-Aldrich or provided by Evonik, as substrate and

the calibration curve was performed with a N-acetyl-D-glucosamine (GlcNAc) solution of 3 mg/mL for the quantification of reducing sugars. One unit (U) of activity corresponded to the release of one  $\mu$ L of reducing sugars per minute.

- Prepare a 10 mL substrate solution by mixing 0.4-10.0 mg mL<sup>-1</sup> hyaluronic acid in 20 mM phosphate buffer solution pH 6.3 (dissolve 2.5 g of monobasic sodium phosphate, 1.0 g of anhydrous dibasic sodium phosphate, and 8.2 g of sodium chloride in water to make 1 L). Any other buffer can be used.
- Pipette 100 μL of enzyme solution into 9.9 mL substrate solution, into a close vial with vigorous mixing.
- Incubate at the desired temperature for 1-24 h.
- Take aliquots and measure the release of reducing sugars as follows.

## 6.1.1. DNS method

The activity of the hyaluronidase is determined by detection of reducing sugars with a modified 3,5dinitrosalicylic (DNS) method adapted to a 96-well microplate scale.

- Prepare the DNS solution: 10 g 3,5-dinitrosalicylic acid (DNS), 300 g sodium and potassium tartrate (Rochelle salt), 16 g NaOH and distilled water up to 1 L (*Note: it takes some time until completely dissolved*).
- In a microtiter plate or 1.5 mL Eppendorf, add 45 μL buffer, 5 μL reaction solution and 50 μL DNS solution.
- Incubate at 85°C for 25 min (in these conditions, the reducing sugars allow the transformation of DNS into DAS with the color change from yellow to dark brown).
- Add 100 μL water and read the absorbance at 540 nm.

### 6.1.2. BSA method

The hyaluronate lyase activity can be additionally determined by quantifying the unsaturated degradation product 2-acetamido-2-deoxy-3-O-( $\beta$ -d-gluco-4-enepyranosyluronic acid)-d-glucosamine ( $\Delta$ Di-HA) photometrically at 232 nm (adapted from Oettl M, Hoechstetter J, Asen I, Bernhardt G, Buschauer A. Comparative characterization of bovine testicular hyaluronidase and a hyaluronate lyase from *Streptococcus agalactiae* in pharmaceutical preparations. Eur J Pharm Sci. 2003 Mar;18(3-4):267-77):

- Prepare incubation mixture containing 300 μL of citrate-phosphate buffer citrate-phosphate (solution A: 200 mM Na<sub>2</sub>HPO<sub>4</sub>/100 mM NaCl, solution B: 100 mM citric acid/100 mM NaCl; solutions A and B are mixed in appropriate portions to adjust the required pH), 150 μL of water, 100 μL of BSA (0.2 mg BSA per mL of water=solution 1) and 250 μL of substrate (0.4-10 mg hyaluronic acid per mL; solution 3).
- After incubation at 37°C for 10 min, monitor the increase in absorbance at 232 nm.
- Calculate the velocity of the enzymatic reaction from the increase in absorbance ( $\Delta A$ ) at 232 nm per time ( $\Delta t$ ) according to the following equation:

$$Rate \ \left(\frac{\mu mol}{\min L}\right) = \frac{\Delta Abs}{\Delta t \ \varepsilon l}$$

where  $\varepsilon$ =4550 L mol<sup>-1</sup> cm<sup>-1</sup>. The molar absorptivity ( $\varepsilon_{232}$ ) of hyaluronate unsaturated disaccharide ( $\Delta$ Di-HA) was determined at 37°C by measuring the absorbance of a known concentration of  $\Delta$ Di-HA (sodium salt, Calbiochem<sup>\*</sup>, Bad Soden, Germany) at 232 nm under the conditions prevalent in the incubation mixture.

### 6.1.3. BCA method

The bicinchoninic acid (BCA) method is mainly used in protein quantification, but has also been adapted to measure reducing sugars. The reaction with reducing agents results in the reduction of Cu<sup>2+</sup> to Cu<sup>+</sup>, followed by the chelation of one Cu<sup>+</sup> with two BCA molecules, forming an intense purple color with an absorbance maximum at 562 nm. See Bicinchoninic Acid Kit for Protein Determination available from Merck/Sigma.

#### 6.1.4. DMAB method

This colorimetric method is based on the Morgan-Elson reaction modified by Reising et al (see Takahashi T, Ikegami-Kawai M, Okuda R, Suzuki K. A fluorimetric Morgan-Elson assay method for hyaluronidase activity. Anal Biochem. 2003 Nov 15;322(2):257-63; J. Biol. Chem., 217 (1955), pp. 959-966). It is based upon the generation of a new reducing GlcNAc terminus with each cleavage reaction. Briefly, after reactions:

- Take 200 µL of reaction mixture.
- The Morgan–Elson color reaction was started by the addition of 50 μL of tetraborate reagent<sup>1</sup> and subsequent heating for 3 min in a boiling water bath.
- After cooling to room temperature, add 1.5 ml of p-Dimethylaminobenzaldehyde (DMAB) reagent<sup>1</sup> and incubate it at 37°C for 15 min.
- After centrifugation at 18,000 g at 4°C for 10 min to remove turbidity, measure the absorbance at 585 nm (or 450 and 650 nm) of the clear supernatant against that of a blank test, which was carried out in the absence of enzyme. Alternatively, the released reducing terminal GlcNAc in the supernatant can be detected by fluorescence (excitation, 545 nm; emission, 604 nm) in a fluorescence spectrophotometer, instead of absorbance at 585 nm.

In both the colorimetric and the fluorimetric methods, 1 unit of hyaluronidase activity is defined as the amount of enzyme required to produce 1  $\mu$ mol of reducing terminal GlcNAc per minute under the specified conditions, using respective standard curves plotted with known concentrations of GlcNAc.

<sup>1</sup>The Reissig method needs two solutions (Analytical Biochemistry293,53–59 (2001)): (i) the borate solution prepared by dissolving  $K_2B_4O_7 \cdot 4H_2O$  at 0.8 M in Milli-Q water (Waters) and (ii) 0.1 g mL<sup>-1</sup> DMAB solution prepared by dissolving 5 g DMAB (Sigma D8904) in 6.25 mL hydrochloric acid, 12 N (Sigma H 7020), made up to a final volume of 50 mL with glacial acetic acid (Sigma A 6283). The latter solution was diluted 10-fold with glacial acetic acid just before use (and at least 15 min before use).

## 6.2. Liquid analysis of hyaluronic acid oligosaccharides by HPAEC-PAD

Commercial hyaluronic acid oligosaccharides can be used: Hyalo-Oligo, a low molecular weight hyaluronan oligosaccharide mixture with an average molecular weight below 10 KDa. It was kindly provided by Kewpie Corp. The mixture is analyzed by high performance anion exchange chromatography with pulsed amperometric detection (HPAEC-PAD) with a Dionex ICS3000. An anion exchange carbo-pack PA 100 column connected to a PA100 guard column is used at 30°C.


## 6.3. Liquid analysis of hyaluronic acid oligosaccharides by SEC-ELSD

Hydrolysis of hyaluronic acid is analyzed by high performance size exclusion chromatography with evaporative light scattering detector (HPLC-SEC). We could study the hyaluronic acid consumption and the formation of the hyaluronic acid oligosaccharides (HAOS). A PolySep-P 4000 column (Phenomenex) is used at 30°C. The analysis is performed at isocratic conditions using milliQ H<sub>2</sub>O as mobile phase at 0.6 mL/min. The samples show a wide molar mass distribution. In SEC, the molar mass of the polysaccharides is estimated from a calibration curve, which is normally obtained from well-defined standards. However, we do not have different size HAOS standards.

## 6.4. Agar assay for the determination of hyaluronidase activity

The agar assay method is adapted from Patil & Chaudhari 2016 (J Basic Microbiol. 2017; 57: 358–361).

- Add hyaluronic acid (0.4-10.0 mg mL<sup>-1</sup>) as a substrate and agarose/agar (1%) as a solidifying agent into the medium where microbes grow or into the buffer where enzymes function.
- Sterilize the medium/buffer i and pour it into Petri dishes up to a depth of 3-4 mm.
- Then, add 50 μL of 24 h old culture supernatant of microorganisms, or 50 μL of enzyme solution under investigation.
- Incubate the plates at 37°C (or any other temperature) for 1-24 h.
- Upon incubation, flood plates with Gram's iodine for 1 min in order to visualize the zones of clearance around the wells (see image below).



## Annex - List of esters successfully tested with pH shift assay


Name	Chemical structure
(1R)-(-)-Dimenthyl succinate	
1,10-Diacetoxydecane	
1-Methyl L-Glutamate	→ → → → → → → → → → → → → → → → → → →
1-Naphthyl butyrate	
1-Napthyl acetate	
2-(tert-Butylamino)ethyl methacrylate	
2,2,3,3,4,4,5,5-Octafluoro-1,6- hexyl diacrylate	O FF FF O FF FF FF FF O
2,2,3,3,4,4,5,5-Octafluoropentyl acrylate	
2,3,4,9-Tetrahydro-1h-b- carboline-3-carboxylic acid methyl ester	H NH O O
2,4-Dichlorobenzyl 2,4- dichlorobenzoate	


2,4-Dichlorophenyl 2,4- dichlorobenzoate	
2-Naphthyl acrylate	
3-beta-Hydroxy-16-alpha- methyl-5-pregnen-20-one 3- acetate	
3-Formylphenyl 3,5- dimethoxybenzoate	
3-Iodo-1H-indazole-6-carboxylic acid methyl ester	
4,6-Dimethyl-α-pyron	
4-Glycylphenyl benzoate	NH <sub>2</sub>
6-Chloro-8-methyl-imidazo[1,2- a]pyridine-2-carboxylic acid ethyl ester	
Allyl cyclohexanepropionate	
Benzyl (R)-(+)-2-hydroxy-3- phenylpropionate	
Benzylparaben	HO

Bis(2-hydroxyethyl) terephthalate	но от
Butyl acetate	$\sim$
Butylparaben	HO
Citraconic acid dimethyl ester	
Cyclandelate	OH OH
Cyclohexyl butyrate	
Dibenzyl terephthalate	
Diethyl adipate	
Diethyl ethoxymethylenemalonate	
Diethyl furan-2,5-dicarboxylate	
Diethyl sebacate	
Diethyl succinate	

Diethyl-2,6-dimethyl 4-phenyl- 1,4-dihydro pyridine-3,5- dicarboxylate	
Dimethyl 1,3-thiazolidine-2,4- dicarboxylate	
Dimethyl 5-(2-amino-4- (methoxycarbonyl) phenoxy)isophthalate	NH <sub>2</sub>
Dimethyl terephthalate	
Dodecanoyl acetate	$\sim\sim\sim\sim\sim\sim$
Ethyl 2-(5-amino-3-tert-butyl-1H- pyrazol-1-yl)acetate	$\rightarrow$ $N^{-N}$ $O^{-N}$
Ethyl 2,5-dibromofuran-3- carboxylate	
Ethyl 2-chlorobenzoate	
Ethyl 2-ethylacetoacetate	
Ethyl 2-methylacetoacetate	
Ethyl 3-(4-methoxyphenyl)-1- methyl-1H-pyrazole-5- carboxylate	

Ethyl 3,5-dimethyl-1H-pyrazole- 4-carboxylate	N. HN
Ethyl 3-Aminopyridazine-4- carboxylate	N NH2
Ethyl 3-oxohexanoate	$\sim$
Ethyl 4-(3,4- dimethoxybenzylamino)benzoat e	
Ethyl 4,4,4-trifluoro-2-butynoate	F F F F
Ethyl 6-aminopyridazine-3- carboxylate	$H_2N$ $H_2N$ $O$
Ethyl acetate	$\mathcal{A}_{0}$
Ethyl acetoacetate	
Ethyl benzoate	
Ethyl beta-morpholinocrotonate	
Ethyl butyrate	
Ethyl decanoate	$\sim \sim $
Ethyl dodecanoate	$\sim\sim\sim\sim\sim$
Ethyl hexanoate	$\sim \sim $





	O Ö
Methyl 2-methoxy-5- (methylsulphonyl)benzoate	
Methyl 3-amino-5-bromo-2- methylbenzoate	Br O NH <sub>2</sub>
Methyl 3-hydroxybenzoate	HO
Methyl 4-bromo-1,2,5- thiadiazole-3-carboxylate	
Methyl 4-hydroxy-1H-indole-6- carboxylate	
Methyl 5-bromo-4- chlorosalicylate	Br CI OH
Methyl 5-fluoronicotinate	FO
Methyl 5-methylhex-2-enoate	
Methyl 6-bromo-7-chloro-8- methyl-4-hydroxyquinoline-2- carboxylate	
Methyl benzoate	
Methyl butyrate	
Methyl cinnamate	

Methyl decanoate	$\sim$
Methyl dodecanoate	
Methyl ferulate	
Methyl glycolate	HO
Methyl hexanoate	
Methyl imidazo[1,2-b]pyridazine- 6-carboxylate	
Methyl myristate	$\sim \sim $
Methyl octanoate	
Methyl oleate	
Moguisteine	
n-Pentyl benzoate	
Octyl acetate	$\sim$
Octyl gallate	
Oxybutynin	

Pantolactone	
Pentabromophenyl acrylate	O $Br$ $Br$ $Br$ $Br$ $Br$ $Br$ $Br$ $Br$
Pentadecyl acetate	
Pentane-1,5-diyl diacrylate	
Phenyl acetate	Î.
Phenyl propionate	
Phenylethyl cinnamate	
Phthalic acid diethyl ester	
Phthalic acid diethyl ester	
Polycaprolactone diol	но~о~он
Propyl acetate	
Propyl butyrate	
Propyl hexanoate	
Propyl propionate	
Propylparaben	но

Sec-Butyl methacrylate	$\sim$
Tetracaine	
Tetramethyl 1,2,4,5- benzenetetracarboxylate	
trans-4'-Cyano-[1,1'-biphenyl]-4- yl 4-ethylcyclohexanecarboxylate	
Tri(propylene glycol) diacrylate, mixture of isomers	$ \qquad \qquad$
Vinyl acetate	$\mathcal{A}_{0}$
Vinyl acrylate	
Vinyl benzoate	
Vinyl butyrate	
Vinyl crotonate	
Vinyl laurate	
Vinyl myristate	
Vinyl oleate	
Vinyl palmitate	
Vinyl propionate	

α-Angelicalacton	
γ-Dodecalactone	
δ-Dodecalactone	
ε-Caprolacton	
Y-Valerolactone	$\begin{pmatrix} 0 \\ \end{pmatrix} = 0$